

Sand Hill Lakes Mitigation Bank
Fourth Annual Report
December 2009



Executive Summary

The Sand Hill Lakes Mitigation Bank (SLMB) consists of approximately 2,155 acres in southern Washington Co. in the Sand Hill Lakes region of the Florida Panhandle (Figure 1). It is located just west of the intersection of SR77 and SR 279 within Township 1 North, Range 14 & 15 West. It contains approximately 850 acres of wetlands, 155 acres of natural lakes and ponds, and 1,150 acres of upland buffer communities. The FDEP permit for the SHLMB was issued September 5, 2005. This is the fourth annual report for the SHLMB. A synoptic listing of notable activities conducted prior to this report and those anticipated in the coming year are presented below.

Interim Success Criteria

All interim success criteria have been met since January 2010. A sufficient number of both natural cypress regeneration and black gum seedlings (300 t/ac) planted along the shoreline of the slough area (Dykes Mill Pond shoreline) and have survived at least one year (January 2010). A sufficient number of planted long leaf pine (436/acre) have survived at least one year for the sand hill restoration (formally pine plantation 319 acres) (January 2009). The following interim success criteria have been met since 2007: All hydrologic and erosion restoration activities have been completed, the Dykes Mill pond dam has been removed in slough area; prescribed burns have occurred in accordance with the burn plans; and all erosion areas, road removal, dam replacement at Black pond and culvert replacement has been completed. All natural and planted areas have increasing vegetation, less than 2% exotic vegetation; preservation areas are maintaining or improving function; upland and wet pine flatwoods have measurably increased in herbaceous ground cover and decreasing in woody vegetative cover; adequate numbers of pine exist within the polygons to meet permit requirements. The targeted oaks (Management Unit 12 and portion of Management Unit 10 were reduced in number in 2006/2007. In 2008, the re-sprouting of oaks was steadily increasing. In 2009, it was determined that the oak densities had increased to greater than 300 trees per acre and should be further reduced to maintain an oak density of less than 150/trees per acre. In June of 2009, ULW was applied at a rate of 1.67 lbs per acre to further reduce the oak re-sprouts in Management Unit 12 and portions of Management Unit 10. Oak re-sprouts were significantly reduced by the ULW to significantly fewer than 150 trees per acre. A total of 640 acres were burned in Management Unit 2, 3, 10, 11, 12, was burned during the winter of 2009 and 2010.

Restoration Activities Completed

Perimeter fencing, gates and signage were installed by February 2005. Ongoing law enforcement has been conducted at the site since 2003 with the purchase of the bank property with no violations to date. A conservation easement was recorded for the SHLMB in February of 2006, preserving the wetland, aquatic and upland communities in perpetuity. Duncan Cairns, Tyler Macmillan and David Clayton were approved as QMS officers for the SHLMB. In accordance with permit requirements a mitigation fund was established for the bank. An archeological/historic survey was conducted at the SHLMB and approved by the Florida Division of Historical Resources (DHR). The majority of the restoration activities were to be initiated during 2005/2006. However, due in part to the delayed permit approvals and a lengthy archeological review by DHR, the initiation of many mitigation activities were initiated approximately a year from the proposed timeline. The initial replanting long leaf pine in the uplands surrounding Black Pond, Cat Pond, and Greenhead branch occurred in December 2004. Oak reduction in the uplands was initiated in June 2005 and completed in August 2006. Construction activities were initiated in July of 2006, in accordance with all permit requirements. All stabilization of erosion areas and re-vegetation, road fill removal, bridge and culvert replacement, Dykes Mill Pond dam removal were completed by March, 2007. The replacement of the dam at Black pond was initiated in October 2007 and was completed by January 2008. Removal of off- site sand pine and slash pine plantation was initiated in July 2007 and completed by October 2007. Gyro-Trac work was initiated in March of 2007 and completed by September of 2007. An additional 38 acres Gyro-Trac work was conducted in June and August 2008 in areas that were historically wet pine flatwoods bringing the total acreage of wet pine flatwood

restoration to 165 acres, an increase from the original 147 acres. Long leaf pine planting occurred in the uplands during the winters of 2005 and 2007. Additional long leaf pine planting occurred in November 2008 in Management Unit 11 where the offsite sand pine or slash pine had been harvested (319 acres). Pond cypress was planted at Dykes Mill Pond in January 2008 and during the drought a significant number of natural cypress seedlings were observed. In addition, a supplemental planting of black gum occurred in May of 2009. Initial wire grass plantings occurred in the wet pine flatwoods in 2006. Wire grass planting is ongoing as allowed in the permit due to the large acreage to restore and limited seed source. To date a total of 165 acres of wet pine flatwoods/wet prairie habitat has been replanted in wet wire grass tublings on 3' centers for a total of 798,600 plugs. Similarly, a total of 110 acres of upland wire grass tublings have been planted for a total of 532,400 tublings. An additional 30 acres of upland and wetland wire grass tublings will be planted in winter 2010/2011.

Controlled Burns

Fire was re-introduced to the SHLMB in the fall of 2004. All initial burns for the wetlands and uplands were completed in December 2006. A total of 2 burn cycles have been completed for the entire SHLMB to date although areas requiring more frequent fires such as sand hills and wet flatwoods have had as many as 4 burn cycles since the bank was established. In 2007, summer burns were re-introduced to portions of the bank. Warm season burns will be conducted in 2008 for 368 acres. A total of 640 acres were burned in Management Unit 2, 3, 10, 11, 12, during the winter of 2009 and 2010. Winter burns were needed due to the higher fuel loads generated from herbiciding the oaks.

Nuisance and Exotic Species

Surveys of nuisance species (flora and fauna) have been conducted throughout the past 5 years. In addition a yearly fall site inspection for nuisance species occurs in conjunction with the annual monitoring as well as day to day monitoring by District and FWC staff. Several small patches of torpedo grass (*Panicum repens*) were treated with Habitat at historic boat launch areas during August and September 2005 and 2007. No live plants were observed during the fall monitoring in 2009. Minor feral hog damage was observed at Dry and Dykes Mill Pond in 2007. Very limited signs of hogs were observed in 2008 and 2009. Water level gages were installed and surveyed in on December of 2005 for 10 locations throughout the bank, and have been read by the FWC for the last three years and data supplied to the District.

Annual Sampling

The annual sampling for this report was conducted in October 30- November 5, 2009. Pedestrian surveys were conducted for both wetland and uplands. The pedestrian surveys were very useful in providing detailed species lists and a greater understanding of species diversity for each community. In addition the pedestrian surveys cover far more area of the polygon that may reveal late successional and threatened or endangered species. In Pedestrian surveys are also useful in identifying pockets of nuisance species and determine fuel loads. Overall, species diversity was excellent throughout the SHLMB and plants were healthy. The number of species observed continues to increase as habitats improve.

Table of Contents

Executive summary	2-3
Table of contents	4
List of figures and tables	5
Introduction	6
Bank establishment and implementation	8
Work schedule	8 - 9
Hydrologic enhancements	9 - 11
Fire management	12-15
Exotic fauna and vegetation	16 -17
Monthly water gage assessments	17 - 20
Sand hill restoration	20 - 25
Wet pine flatwood restoration	26 - 28
Annual monitoring	28 - 29
Quantitative monitoring	29 - 48
Qualitative monitoring	48 - 56
Appendix 3: Computations and analysis	http://www.nfwmdwetlands.com
Appendix 4: Pedestrian Surveys and transect photography	http://www.nfwmdwetlands.com
Appendix 5: Aerial oblique photography	http://www.nfwmdwetlands.com
Appendix 6: Florida Fish and Wildlife Conservation Commission	http://www.nfwmdwetlands.com

List of Figures and Tables

Figure 1.	Location map	6
Figure 1a.	SHLMB boundary map and habitat	7
Figure 2.	Road improvements	9
Figure 3.	Structures	10
Figure 4.	Erosion and Stabilization Sites	11
Figure 5.	Anticipated Burn Cycles	13
Figure 6.	Areas Burned Since inception of the Bank through 2006	14
Figure 7.	SHLMB 2009 Burns	15
Figure 8.	Nuisance and Exotic Species Tracking	17
Figure 9.	Water Level Staff Gage Locations	19
Figure 10.	Oak Removed Through 2006	21
Figure 10a	2009 Oak Eradication	22
Figure 11.	Management Unit 11 Pine Removal	23
Figure 11a.	Sandhill Long Leaf Pine Planting	24
Figure 11b.	Wet Prairie Wire Grass Planting Location	25
Figure 12.	Pine Flatwoods Restoration Areas	27
Figure 12a.	Wet Prairie Wire Grass Planting Area (2009)	28
Figure 13.	Monitoring Locations	30
Figure 14.	Transect 1: Species cover and occurrence (Sand Pine Plantation)	32
Figure 15.	Transect 2: Species cover and occurrence (Sand Pine Plantation)	34
Figure 16.	Transect 4: Species cover and occurrence (Sand Pine Plantation)	35
Figure 17.	Transect 3: Species cover and occurrence (Sand Hill)	38
Figure 18.	Transect 5; Species cover and occurrence (Sand Hill)	39
Figure 19.	Transect 8: Species cover and occurrence (Slash Pine Plantation)	42
Figure 20.	Transect 6: Species cover and occurrence (Hydric Pine Flatwoods)	45
Figure 21.	Transect 7: Species cover and occurrence (Hydric Pine Flatwoods)	46
Figure 22.	Transect 9. Species and occurrence (Slough)	48

Tables

Table 1.	Revised work schedule	9
Table 2.	Water level staff gage readings - 2007	18
Table 3.	Transect 1. Species cover and occurrence (Sand Pine Plantation)	31
Table 4.	Transect 2. Species cover and occurrence (Sand Pine Plantation)	32
Table 5.	Transect 4. Species cover and occurrence (Sand Pine Plantation)	34
Table 6.	Transect 3. Species cover and occurrence (Sand Hill)	37
Table 7.	Transect 5. Species and occurrence (Sand Hill)	38
Table 8.	Transect 8. Species and occurrence (Hydric Pine)	41
Table 9.	Transect 6. Species and occurrence (Hydric Pine)	44
Table 10.	Transect 7. Species and occurrence (Hydric Pine)	45
Table 11.	Transect 9. Species and occurrence (Slough / Marsh)	47

Introduction

The Sand Hill Lakes Mitigation Bank (SLMB) consists of 2,155 acres in the southern portion of Washington Co. in the Sand Hill Lakes region of the Florida Panhandle (Figure 1). It is located just west of the intersection

of SR77 and SR 279, and is within Township 1 North, Range 14 & 15 West. It contains approximately 850 acres of wetlands including high quality cypress sloughs and strands, degraded hydric pine flatwoods, bayheads, seepage slopes, and approximately 155 acres of natural solution ponds and shallow, gently-sloped lakes connected by streams and ditches. The remaining 1,150 acres consist of secondary growth upland buffer communities (including high quality and degraded sand hill communities as well as sand pine plantation, slash pine plantation, and mixed hardwoods) (Figure 1a).

The SHLMB occurs on the divide between the Choctawhatchee and St. Andrew Bay watersheds. The majority of the proposed Bank is in the surface headwaters of Pine Log Creek, which flows westerly and southwesterly to Pine Log State Forest and ultimately to the Choctawhatchee River and Bay. However, because of the karst nature of the Sand Hill Lakes the SHLMB is also a recharge area for Econfina Creek, which, via Deer Point Lake, is the water supply for Panama City.






Figure 1a. SHLMB boundary map and habitats

Sand Hill Lakes Mitigation Bank - 1999 DOQ



Bank Establishment and Implementation of Permit Requirements

The permit for the Sand Hill Lakes Mitigation Bank (SHLMB) was issued by the DEP on September 5, 2005. This document represents the fourth annual report for the SHLMB. Perimeter fencing with gates and signs were installed prior to March of 2005. Law enforcement has been conducted at the site since the property was purchased and is ongoing at the SHLMB. A conservation easement was recorded for the SHLMB on 2/28/06, preserving the wetland, aquatic and upland communities in perpetuity. QMS officers Duncan Cairns, Tyler Macmillan and David Clayton were selected by the NFWFMD and approved by the DEP. In accordance with permit requirements a mitigation fund was established for the bank. An archeological and historic survey was conducted for the SHLMB and approved by the Division of Historical Resources. Construction activities were initiated in July of 2006, in accordance with permit requirements.

Mitigation Activities Work Schedule

According to the proposed work schedule for the SHLMB found on page 12 of the SHLMB permit, the majority of the restoration activities were to be initiated during 2005-2006. However, the restoration activities were postponed due to delays in permit issuance, recording of conservation, and additional time needed to complete and approve the archeological study. Consequently, many of the restoration activities were delayed by approximately 1 year. A revised schedule was included in the first monitoring report. In 2008, the majority of the construction and restoration activities were completed and an updated work schedule has been provided (Table 1).

Table 1. Restoration work schedule

Activity	Estimated Completion Date
Conservation easement, QMS	Completed 3/06
Fencing and signage	Completed 3/05
Site security / law enforcement / internal gating / road closures	Ongoing
Stabilization of 10 erosion sites	Completed 3/2007
Hydrologic enhancements - Replacement of Black Pond dam - Removal of Dykes Mill Pond dam - Removal of road fill at (3) sites - Construction of 2 bridges and replacement of 3 culverts	Initiated 10/07 Completed 1/08 Initiated 7/06 Completed 8/06 Initiated 7/06 Completed 3/07 Initiated 7/06 Completed 3/07
Removal of pine plantation and thinning of slash pine	Initiated 7/07 Completed 10/2007
Removal of oak overgrowth and replanting with longleaf pine	Completed: Oak removed 2005/2006, additional oak removal (ULW, 6/2008) Pine planted 2005 and 12/2007
80% completion of initial growing season and fuel reduction fires in areas to be maintained as oak / pine community	Completed 12/2005
Initial thinning, roller chopping, and fuel reduction fires in hydric pine	Completed Initial burns 8/05 Completed required shrub reduction 6/07 (Gyrotrack) Completed Pine thinning 10/07 Completed site prep burns following harvest 12/2008
Supplemental wiregrass seeding if necessitated by onsite conditions	2008/2012 Ongoing Initial wet flatwoods wire grass planting (165 acres) completed 12/09
Installation of water level gages	Completed 12/05
Baseline assessments of vegetation	Completed 2004/2005
Fire Management / Monitoring Year 1 / Annual Report	Completed 2005/2006 report

Fire Management / Monitoring Year 2/ Annual Report	Completed 2007/2008 report,
Fire Management / Monitoring Year3 / Annual Report	Completed 2009/2010 report
Fire Management / Monitoring Year 4 / Annual Report	2010/2011 report
Fire Management / Monitoring Year 5 / Annual Report	2012/2013 report
Perpetual Ecological Management	2013 +

Hydrologic Enhancements

Hydrologic enhancements include the complete removal of 2 fill-road crossings, installation of bridges at 3 crossings and 2 culverts and the removal or replacement of 2 failing water control structures, the remediation of 10 erosion areas, the stabilization of 1 boat launching site, and construction of one rain shelter (Figures 3 and 4).

The removal of the failing water control structure at Dykes Mill Pond and construction of three bridges (#1, #3, #7), and two culverts (#9, #10-A-B) was initiated in July 2006 and completed in April of 2007 in accordance with permit conditions (Figure 3). The graded areas were stabilized and seeded in early 2007 with season-appropriate, non-invasive annual grass to reduce potentially turbid runoff. On June 30th, the graded areas were seeded with brown-top millet. Currently all water control structures are functioning properly and water levels have risen to post drought levels.

The removal and re-vegetation of two fill-road crossing was initiated in January of 2007 and completed in March of 2007 (Figure 3). Erosion area #6 was restored in July of 2006 as part of road enhancement project while remediation of the remaining 9 erosion sites was initiated in January and completed in April 2007. Hay bales and silt fences were installed in accordance with the permit requirements (Figure 3 and 4). The areas were planted as each site was completed. Sites were planted in accordance with the approved planting plan. Graded areas were stabilized with annual rye grass and seeded with brown-top millet on June 30, 2007. Sites were monitored during the summer and fall monitoring. Inadvertently, the contractor used Bahia grass hay to stabilize soils at the two erosion areas 1-3 (Cat Pond and the road removal at Deep Edge). The contractor was required to treat each area with herbicide until the Bahia grass was eliminated. Initial treatments occurred in May with subsequent treatments in September. Supplemental wire grass and long leaf pine seedlings will be planted at these sites in 2008. In 2008 seed from the eradicated Bahia grass and or mulch germinated as was again reated with to remove the grass. In addition, poor survival was observed at the erosion sites 1, 2, 4, 5, and 10. Supplemental planting occurred in February 2008 in accordance with the permit requirements. Shrubs were planted at the road fill removal sites in March 2009 in accordance with the planting plan. This action completes the planting requirements for these areas.

The replacement of the water control structure at Black Pond (#2) was initiated in October 31, 2007 and completed by the end of January 2008.

Finally, the stabilization of one boat launch area on Dry Pond was completed in September 2007. Photographic documentation for all these activities was included in the 2007 report.

Figure 3 - Structures





Figure 4 - Erosion Stabilization Sites



0 0.5 1 Miles



Fire Management

The bank is divided into 14 Management Units that range from 0.25 to ~580 acres. Prescribed fire is an integral component of the management, enhancement and restoration for six of the management units (Management Units 2, 3, 8, 10, 11, and 12), and will also be used to manage portions of the power line ROW (Figure 5). The remaining Management Units are wetlands or aquatic systems not typically managed with fire, although fire from adjacent Units may be allowed to burn into them when conditions allow. Prescribed burns have generally been conducted during the growing season (March through August), although initial dormant-season fuel-reduction fires have been required in some areas. Burns are planned for 1-3, 3-5 and 5-7 year cycles, although fuel levels, prevailing weather patterns and other on-site conditions may necessitate modification of burn cycles. Burn coverage of 80% or more within a polygon has been considered a successful burn. Prescribed fire is intended to inhibit establishment of woody species, promote fire-adapted species, and stimulate seed production of desirable herbs. Fire prescriptions have been written to comply with open burning laws (Florida Statutes 590) and liability considerations. Safety and protection of property will be the priority concern of the Florida Certified Prescribed Burn Manager (FCMB).

Fire was re-introduced to the SHLMB during the winter of 2004 to portions of Management Unit 11 and 12. Subsequently portions of the sand hills and hydric pine flatwoods were burned during the summer of 2005 with the remaining initial burns completed by December of 2005 in accordance with the Fire Management Plan (Figure 6). In areas with a high fuel loads such as Management Unit 2, 3, 8 and some portions of Management Unit 10 adjacent to Black pond dormant season fuel reduction fires were utilized. However in Management Unit 11 and 12 initial burns reduced fuel loads to the extent that warm season burns were conducted. Wire grass flowered in these areas following the fire and plants appeared healthy. The burns at the SHLMB have also been successful in reducing woody vegetation coverage as well as stimulating a seed bank of fire adapted species. Prior to the initiation of fire, woody goldenrod was the dominant herbaceous species, but the initial fires greatly reduced the woody goldenrod cover and stimulated the wire grass. Currently wire grass is the dominant herbaceous species and the sand hills and wet flatwoods have greatly reduced shrub cover.

In 2007, it was anticipated that 287 acres would be burned during winter 2006/2007. However, due to the extended drought and unsafe fire conditions only 69 acres were burned with 66 acres meeting the 80% requirement. No warm season burns were attempted due to the extended drought.

In 2008, a total of 384 acres of burns are planned at the SHLMB (Figures 6a, 6b, 7, 7a-7e)). Warm season burns will be conducted at Garret Pond and adjacent to Pine Log Creek and winter burns will be conducted for the areas with off-site pine removal.

As of 2009, a total of 2 burn cycles have been completed for the entire SHLMB to date although areas requiring more frequent fires such as sand hills and wet flatwoods have had as many as 4 burn cycles completed. During the winter of 2009 a total of 600+ acres were burned in Management Unit 2, 3, 10, 11, 12. Hydric flatwood wetland areas were burned in preparation for planting of wire grass or to reduce the dead shrubs that had been treated with herbicide (Table 6a). In the uplands the sand hill areas that had been treated with ULW to reduce oak coverage were also burned to reduce standing dead shrubs and to stimulate the herbaceous layer.

Figure 5 - Anticipated Burn Cycles




Figure 6 - Areas Burned Since Inception of Bank Through 2006






Figure 7. SHLMB 2009 Burns



Exotic Fauna and Vegetation

Surveys nuisance species have been conducted throughout the year. In 2006, one female hog was trapped and patches of torpedo grass were observed at historic boat launches. These areas were treated by the Bureau of Invasive Plant Management on July 20th 2006. In 2007, several small patches of torpedo grass were again observed at the historic boat launch areas of several ponds (Figure 8). These areas were treated twice with Habitat on July 26th and August 6, 2007. No visible living plant material was observed during subsequent site visits or during the 2008 fall monitoring. Inadvertently the contractor working on the road removal and stabilization of erosion areas used inappropriate Bahia grass hay to stabilize the soils for erosion areas 1 and 2 and 3 the road removal between Deep Edge and Little Deep Edge. The contractor was required to treat these areas with herbicide until the Bahia grass was killed. Treatments occurred in May and September. No living material was observed during the fall monitoring. In the summer of 2008, small patches of Bahia grass developed from the seed bank in the road removal areas and were again treated twice in 2009. In addition, scattered Bahia grass plants were observed on 83 acres adjacent to Green Head Branch that had been planted in sand pine and 53 acres adjacent to the check station that had been planted in slash pine. Based on weeds associated with these areas, it is probably that these sites had been used for agriculture or pasture prior to conversion to sand pine plantations. The Bahia grass in these areas was treated by hand crews twice in 2009 and Bahia grass cover was greatly reduced. In 2010 hand treatment will continue in the areas described above. Some hog damage was observed in 2008 during the drought adjacent to Dry Pond and the Green ponds. However, in 2009, no real hog damage was observed. In 2009, several packs of dogs were observed chasing wildlife harassing day visitors. The dogs were aggressive and had threatened visitors and one bit a staff member. The County Sherriff department, and animal control were contacted and helped in the investigation. Traps were set to capture the dogs. It appears the owner has been located and dogs in question are now under control.

Figure 8 - Nuisance and Exotic Species Tracking



Monthly Water Gage Assessments:

Water levels gauges were installed and surveyed in on December of 2005 for 10 locations throughout the bank. These locations include Black Pond, Power Line Pond, Pine Log Creek, Deep Edge Pond, Little Deep Edge Pond, Dykes Mill Pond, Ditch connecting to Pine Log Creek #7, natural channel from Joiner Lake to the Green Pond, Green Ponds, and Dry Lake (Table 2, Figure 9). The gauges were read monthly by the Florida Wildlife Conservation Commission staff and the results submitted to the NFWFMD (Table 2, Figure 9). In 2006, the water levels were above the gages until April, then from May to December then


Table 2. Monthly Water Gage Readings

Readings in Feet	(1) Black Pond	(2) Power Line Pond	(3) Pine Log Creek	(4) Deep Edge Pond	(5) Little Deep Edge Pond	(6) Dykes Mill Pond	(7) Green Ponds Channel	(8) Joiner Lake Canal	(9) Green Ponds	(10) Dry Pond
Date	1/5/2009	1/5/2009	1/5/2009	1/5/2009	1/5/2009	1/5/2009	1/5/2009	1/5/2009	1/5/2009	1/5/2009
Reading	below gauge	no water	1.72	below gauge	2.52	3.38	3.05	no water	below gauge	2.98
Date	2/12/2009	2/12/2009	2/12/2009	2/12/2009	2/12/2009	2/12/2009	2/12/2009	2/12/2009	2/12/2009	2/12/2009
Reading	below gauge	no water	0.3	below gauge	2.15	3.13	below gauge	no water	below gauge	2.60
Date	3/3/2009	3/3/2009	3/3/2009	3/3/2009	3/3/2009	3/3/2009	3/3/2009	3/3/2009	3/3/2009	3/3/2009
Reading	below gauge	no water	1.98	below gauge	2.44	3.30	2.90	no water	below gauge	2.90
Date	4/3/2009	4/3/2009	4/3/2009	4/3/2009	4/3/2009	4/3/2009	4/3/2009	4/3/2009	4/3/2009	4/3/2009
Reading	1.80	below gauge	3.78	below gauge	2.72	4.40	3.52	4.26	3.30	3.92
Date	5/1/2009	5/1/2009	5/1/2009	5/1/2009	5/1/2009	5/1/2009	5/1/2009	5/1/2009	5/1/2009	5/1/2009
Reading	6.60	6.11	4.12	below gauge	2.52	3.40	3.96	4.44	6.34	>6.70 submerged
Date	6/3/2009	6/3/2009	6/3/2009	6/3/2009	6/3/2009	6/3/2009	6/3/2009	6/3/2009	6/3/2009	6/3/2009
Reading	6.37	5.89	3.88	below gauge	2.52	3.42	3.74	4.58	6.13	6.70
Date	7/7/2009	7/7/2009	7/7/2009	7/7/2009	7/7/2009	7/7/2009	7/7/2009	7/7/2009	7/7/2009	7/7/2009
Reading	5.16	5.22	3.45	below gauge	2.53	3.28	2.70	3.18	4.94	5.49
Date	8/5/2009	8/5/2009	8/5/2009	8/5/2009	8/5/2009	8/5/2009	8/5/2009	8/5/2009	8/5/2009	8/5/2009
Reading	4.58	4.6	2.88	below gauge	2.5	3.27	below gauge	2.52	4.31	4.79
Date	9/1/2009	9/1/2009	9/1/2009	9/1/2009	9/1/2009	9/1/2009	9/1/2009	9/1/2009	9/1/2009	9/1/2009
Reading	4.35	4.17	2.45	below gauge	2.50	3.22	1.75	2.39	4.09	4.61
Date	10/1/2009	10/1/2009	10/1/2009	10/1/2009	10/1/2009	10/1/2009	10/1/2009	10/1/2009	10/1/2009	10/1/2009
Reading	4.60	4.29	2.58	below gauge	2.42	3.29	below gauge	3.09	4.30	4.90
Date	11/2/2009	11/2/2009	11/2/2009	11/2/2009	11/2/2009	11/2/2009	11/2/2009	11/2/2009	11/2/2009	11/2/2009
Reading	4.58	4.16	2.45	below gauge	2.44	3.39	2.42	2.85	4.31	4.90
Date	12/1/2009	12/1/2009	12/1/2009	12/1/2009	12/1/2009	12/1/2009	12/1/2009	12/1/2009	12/1/2009	12/1/2009
Reading	4.45	3.89	2.18	below gauge	2.42	3.30	2.38	2.50	4.20	4.75

<Gage = Water level was down slope of staff gage.

DRY = Site is dry. No data -- site was inaccessible or unread

Figure 9 - Water Level Staff Gage Locations



▲ = Staff Gage (Installed 2005)

0 0.5 1 Miles



water levels were below the staff gages for all but Little Deep Edge and Dykes Mill Pond. The drought continued in 2007 and 2008. In 2009, the extended drought subsided and the Green Ponds that had dried down, along with the Power line pond and Pine Log Creek and Joiner canal were once again filled with water and the water levels at Deep Edge, Black and Dry Pond levels returned to pre-drought levels.

Sand Hill Restoration

Activities: oak eradication, planting of pine, wire grass planting

Oak eradication

A total of 1,150 acres longleaf pine / wiregrass community, live oak forest and other buffer habitats occur on the SHLMB. The NFWFMD will provide perpetual ecological management for these habitats. Oak eradication in Management Unit 12 was completed for the majority of the site in August of 2005 with a small remaining portion completed in September of 2006. Turkey and live oaks were reduced to less than 150 trees per acre and stumps were painted with an approved herbicide to reduce stump sprouts. Similarly, oak coverage was reduced for significant acreage in Management Unit 10 in September of 2006. These areas have excellent wire grass cover and a well developed understory of sand hill species. To date a total of 550 acres of sandhills have had the oaks thinned, far exceeding permit requirements (Figure 10).

In 2007, these areas were again burned, but it was noted that the thinned oaks and hardwoods had re-sprouted and oak densities from the sprouts had increased exceeding target densities. These areas were monitored in 2008, and increasing cover of oaks was observed.

In April of 2009, a comprehensive field review of the uplands at the SHLMB was conducted to determine if additional oak reduction was needed. It was determined that oak numbers from re-sprout were significant and threatened to shade out the wire grass and the oak numbers should be further reduced. Two treatment types were utilized, hand application for small areas or areas adjacent to acceptable oak numbers, and aerial application to treat large areas. The prescription was based on previous experience in similar habitats and a rate of 1.67 pounds per acre of Velpar ULW was applied to 546 acres in May of 2009 (Figure 10a). After approximately 3 weeks noticeable results were observed and after 6 months the oak numbers were significantly reduced. Velpar was applied by a licensed applicator and rates were followed accurately, however, some minor impacts were observed to wire grass cover in scattered locations. Symptoms observed included culms browning and decline. This was not observed widely and it is thought that the sand content of the soil, combined with high temperatures may have caused the problem.


Long Leaf Pine Planting

Prior to permit issuance, longleaf pine seedlings were planted in portions of Management Unit 12 in the winter of 2004. However, intense winter burns in early 2007 destroyed most of the planted. Additional plantings of longleaf pine at a rate of 436 trees per acre occurred in Management Unit 12 and portions of Management Unit 10 during the dormant season of 2007/2008.

Pine Plantation Harvest

Restoration activities for the existing sand pine plantation (~385 acres) and slash pine plantations (11.5 acres) were initiated in June, 2007 (Figure 11). The sand pine and slash pine plantations harvest began on June 15 and completed in November 16, 2007. All sand pine and slash pine scheduled for removal was completed in accordance with permit requirements. These areas were burned in the fall of 2008 and 319 acres were replant in the winter of 2008/2009 with long leaf pine (Figure 11a). The planted long leaf pine established well with excellent survival.

Figure 10 - Oak Removed Through 2006



 Oak Removal Areas Through 2006 (550 Acres)



0 0.5 1 Miles

Figure 10a. 2009 Oak Eradication






Figure 11 - Management Unit 11 Pine Removal



 Pine Removal Areas (~400 Acres)




Figure 11A. 2008 Sand hill longleaf pine planting



Upland Wire Grass Planting

The majority of the areas with sand pine plantation had remnant sand hill species in the understory prior to harvest of the sand pine. Once the sand pine was removed, the sand hill seed bank and remnant wire grass greatly increased in cover in response to added light. However, in areas where few sand hill species were observed or areas without wire grass, wire grass tublings have or will be planted. Due to limited seed source a maximum of 30 acres each year will be planted in accordance with the permit requirements. In 2008, 53 acres of historic sand hill were planted with upland wire grass tublings on 3' centers (256,520 plants). In 2009, an additional 27 acres of upland wire grass tublings were planted on 4' centers (130,680 plants) in areas where the sand hill species were absent (Figure 11b). An additional 30 acres of upland wire grass tublings are scheduled for planting in 2010.

Figure 11b. Wire grass plantings at SHLMB



Northwest Florida Water Management District
Sand Hill Lakes Mitigation Bank (SHLMB)
Section 6, Township 1 North, Range 14 West
Washington Co., Florida



0 0.25 0.5 Miles

Wet Flatwoods Restoration

According to the permit requirements, 147 acres of wet flatwood restoration was scheduled to occur at the SHLMB., Management Unit 2. However, District staff identified and additional 18 acres that historically was wet flatwoods and added this acreage to Management Unit two for a total acreage of 165 acres of wet , flatwoods restoration (Figure 12). Standing biomass of shrubs (primarily titi, gallberry and fetterbush) has been reduced to ground level with the use of a Gyro-Trac followed by winter burns. The gyrotack work was initiated on March 13 and was completed by August 20, 2007. The black titi in these areas was extremely thick often with a dbh of 10-14" and 25 – 30' tall. Even with the large "tree" size black titi, the gyrotrack was excellent in reducing the thick dense shrub cover to ground level. There were no noticeable track marks or ruts left by the Gyro-Track. The mulch within these areas was allowed to dry for several months prior to burning. Sites were burned in December of 2007.


By March 2008, it was apparent, that while the shrub cover was greatly reduced, re-sprouting of the shrubs had occurred in all Gyro-Trac areas. Average shrub densities were determined through randomly established transects and stems per meter squared were determined. In areas with a hot fire 50-80 stems per meter squared were observed while in areas with an incomplete burn, 100 to 135 stems per meter squared were common. Based on these observations, the shrubs would return if not significantly reduced further. In an effort to determine if selected herbicides could aid in reducing shrubs numbers to an acceptable level, two polygons, the Whale, a 12 acre polygon adjacent to the Dry Pond parking, and a 16.2 acre polygon adjacent to Dry Pond and the slash pine restoration sites were chosen. These two areas were treated twice by Entrix with appropriate wetland approved herbicides, once in July and again in September. The 16.2 acre polygon was burned in the winter of 2008-2009 with a very hot fire. The 16.2 acre polygon was planted with wire grass plugs while the whale, had previously been direct seeded with wire grass seed. Preliminary results indicate that the shrub cover was greatly reduced from greater than 85% cover to less than 15% cover. Based on the positive results in these areas, herbicide was used to reduce shrub cover throughout the 165 acres of wet flatwood restoration.

In Management Unit 3, Planted slash pine area restored to wet flatwoods), the shrub layer was limited due to the dense overstory of planted pine. Wet flatwood herbaceous species were more common in these areas after the initial warm season burn (2006) and recent slash pine thinning (2007). The initial fire in this area reduced most of the shrubs to coppice sprouts. It was hoped that the shrubs in this area could be managed through successive warm season fires. The fire was conducted for Management Unit 3 in December 2008. The burn had good coverage and shrubs were reduced to the ground level. However, shrub cover increased following the thinning of the pine, and the area was burned during the winter of 2009/2010. In 2010, the area will be treated with herbicide to further reduce the shrub cover and it will be planted with wet prairie wire grass in the winter of 2010/2011.

Wire grass planting continued in the wet flatwood restoration areas in 2008. A total of 32 acres of wet wire grass tublings (143,880 plants) were planted in December 2008. Tublings were planted on 3 foot centers for the 5.4 acre area north of Dry Pond, the 16.2 acre polygon adjacent to the southern portion of Dry Pond and east of black pond, and the 12.6 acre area surrounding Garret Pond (Figure 12a).

In 2009, the original 165 acres of wet pine flatwoods areas that had shrub reduction were treated with selective herbicides to further reduce shrub cover. Shrubs were hand treated three times in 2009, (April, July and October) to reduce shrub cover without impacting the understory or planted wire grass. This treatment will continue in 2010.

Figure 12. Pine Flatwood Restoration Areas
Brush Reduction



Northwest Florida Water Management District
Sand Hill Lakes Mitigation Bank (SHLMB)
Brush Reduction (Gyro-Track Mulching) - ~165 Acres
Section 6, Township 1 North, Range 14 West
Washington Co., Florida





Figure 12a. Wet wire grass planting areas (Winter 2009).



Annual Monitoring

In accordance with Specific Condition 26, all sampling locations have been identified (Figure 13). Fall monitoring methods as well as data analysis are described below. Raw data, computational analysis, pedestrian surveys and photographic documentation are included in Appendix 2, 3 and 4 and found at the

District website: <http://www.nwfwmdwetlands.com/index.php>. Similarly, Oblique aerials of the SHLMB on October 19, 2009 and can be found at the District website (see above).

The 2008-2009 Annual report by the Florida Fish and Conservation Commission was completed in October and can be found on the District website (see above) in accordance with Specific Condition 25f.

Quantitative Monitoring

Materials and Methods

Quantitative monitoring has been conducted in accordance with the methods described in Attachment H – Monitoring Plan. Quantitative vegetation monitoring occurred at the end of the growing season. This is the second annual monitoring report for the SHLMB.

The percent vegetation cover was monitored at transect locations shown in Figure 13. One-meter square quadrats were established along 600' transects at 20' intervals. In addition, each transect contained a permanently established photographic documentation stations, where qualitative quadrat (north, east, south, and west) observations were recorded (Appendix 4). Transect termini will be marked using iron rebar surrounded by PVC pipe.

Vegetation species coverage statistics were developed from the recorded coverage of each species (or bare ground or open water) within a given quadrat. The percent coverage for each species (and bare ground or open water) was generated by adding all quadrat observations together, and dividing the total coverage by the cover of each species within each transect. This represents a modified Daubenmire cover scale where vegetation species statistics are used to determine the percent cover by bare ground, water, individual species and groups, such as wetland species, invasive exotic and nuisance species, and present.

Tree density was monitored using the “line strip” (belt transect) technique. Transects were co-located with each vegetation transect. The belt transects will be 600± feet in length and 30± feet in width. Within each belt transect, the height and condition of each planted tree will be recorded.

Photographic Stations:


Panoramic photographs were taken from the permanently established stations at each transect and are found on the District website: <http://www.nwfwmdwetlands.com/index.php>. **Please note: photographic station 12 was abandoned as it was not placed in the correct habitat. The photographic station was inadvertently placed in a mesic hammock on the edge of management unit 2. To remedy this, a new photopoint 12b was established in the in management unit 2 to the south of the original photo point (Figure 13).**

Wildlife Utilization:

During the vegetation monitoring described above, wildlife observations will be recorded in each community. These observations will consist of direct sightings, scat, tracks, or vocalizations.

Fuel loads and prescribed fires within wet flatwood and sandhill communities: Semi-annual status reports will detail the condition of the communities relative to the need and potential for a burn, the conditions required for the next desirable burn, and the anticipated timeframe for the next burn. This data was included for each pedestrian survey transect found on the District website (see above).

Figure 13 - Monitoring Locations



Results and Discussion

UMAM Polygon II, Management Unit 11- Sand Pine Plantation

UMAM Polygon II, Management Unit 11, consists of 383.484 acres of planted sand pine plantation that will be converted to long leaf pine and sand hill habitat. Baseline conditions indicated a sand pine canopy with nearly 100 percent canopy closure and an average of 446 sand pine trees per acre occur in the sand pine plantations. Removal of the sand pine was completed in November 2007. Three transects (transect #1, #2 and #4) were located within UMAM Polygon II, Management Unit 11.

In 2008, a total of 10 species were observed in transect 1, 16 in transect 2, and 20 in transect 4. Two transects (1 and 4 lost one and two species respectively, while transect two increased by 11 species (Tables 4-6) (Figures 13-15). Wire grass was observed only in transect 2 with 8.5% cover, an increase of 3% cover from last year and was the dominant species occurring in that transect. The dominant cover class for all transects was bare ground with a range of 82% bare ground (transect 2) to 40% bare ground (transect 4). Bare ground was greatly reduced from the previous year along each transect. The exotic species Bahia grass (*Paspalum notatum*) was observed in transects 1 and increased from 0.1% cover to 0.7% cover. Bahia grass was also observed in transect 4 reduced from 1.5% cover to one percent cover. However, centipede grass increased in cover from 10.6% to

23.2% cover and again was the dominant species within that transect. Herbicide treatments targeting Bahia and centipede grass without impacting the native species will be applied in spring and fall of 2009.

In 2009, a total of 19 species were observed in transect 1, 23 in transect 2, and 23 in transect 4. Increasing species numbers were observed in all transects ranging from an increase of 9 species in transect 1 to 3 in transect 4 (Table 3,4 and Figures 15 and 16). A total of 11 species common to sandhills were found in transect 1, 21 in transect 2, and 17 in transect 4. Since the sand pine plantation was removed there has been increasing numbers of sand hill species present within the transects. Wire grass was observed in transects 2 and 4. Percent cover of wire grass had increased from 8.5% to 25.7% cover in transect 1 and from 3% cover to 12.3 % cover in transect 4 within the last year. Transect 1 will be planted in wire grass tublings during the winter of 2009. Vegetative cover continues to increase for transect 1 and 2 with 35.9 % cover for transect 1, and 34% cover in transect 2. Cover was slightly reduced from 48% cover to 40% cover in transect 4 potentially due to herbicide treatment of the centipede and Bahia grass. Bahia grass was again observed in transect 1 and cover increased from 0.7% cover to 3% cover. Bahia grass was treated last year and will continue to be treated in the following year. However, while Bahia grass cover was greater than desired in transect 1, the cover of Bahia grass is spotty and below 2% cover for the polygon.

Interim Success Criteria:

The sand pine plantation was harvested in 2007. Site preparation burns occurred during the winter of 2008 and the area that included transect 4 and transect 2 was planted in the winter of 2008/2009 with long leaf pine. Wire grass tublings were planted on 3' centers in the polygons surrounding transect 1, and 4 in 2008 and the polygon including transect 1 (22 acres) will be planted in the winter of 2009/2010 (Table 11b). The transects are measurably increasing in species number and vegetation cover, Bahia grass was eliminated from transect 2 and 4, and fire has been re-introduced to the areas.

Table 3. Transect 1 Species cover and occurrence (Former Sand Pine Plantation)

Date 11/4/09
 Collector: David Clayton
 Wildlife observed: Red wing black bird, titmouse, chipping sparrow
 titmouse, chipping sparrow
 Community description: Former Sand Pine Plantation
 Replanted with LLP

Time: 11:00
 Am
 Condition, Fair and cool

<u>Scientific Name</u>	<u>Species</u>	<u>Percent Cover</u>
Aster pilosus	Frost aster	0.5
Artemisia campestris	Wormwood	11.6
Axonopus furcatus	Big carpet grass	0.6
Chrysoma pauciflosculosa	Woody goldenrod	0.5
Cyperus sp.	Sedge	7.6
Dichanthelium aciculare	Needle leaf witch grass	0.66
Diospyros ebenum	Persimon	0.16
Eupatorium capillifolium	Dog fennel	2
Eupatorium mohrii	Mohr's thorough wort	0.5
Mollugo verticillata	Indian chickweed	1
Opuntia humifusa	Prickly pear cactus	0.167

Paspalum notatum	Bahia grass	3.1
Pinus clausa	Sand pine	2.3
Pinus palustris	Long leaf pine	0.83
Quercus hemisphearica	Diamond oak	0.167
Rubus cuneifolius	Sand black berry	1.7
Sida rhombifolia	Indian hemp	0.167
Tradescantia hirsutiflora	Hairy spiderwort	0.33
Yucca filamentosa	Adam's needle	1.6
Bare ground	Bare ground	64.1

Figure 14. Transect 1. Percent Cover and occurrence (Sand Pine Plantation)




Table 4. Transect 2. Species cover and occurrence (Sand Pine Plantation)

Date 11/4/09	Time: 11:30 Am
Collector: David Clayton	Condition, Fair and cool
Wildlife observed: Titmouse, chipping sparrow	Fuel load: Low
titmouse, chipping sparrow	State Threatened species
Community description: Former Sand Pine Plantation	Wire grass and sandhill species regenerating

<u>Scientific Name</u>	<u>Species</u>	<u>Percent Cover</u>
Andropogon arctatus	Pinewoods bluestem	1
Andropogon glomeratus	Bushy broom sedge	0.17
Andropogon virginicus	Broom sedge	0.33
Aristida stricta	Wire grass	25.7
Axonopus furcatus	Big carpet grass	0.16
Cyperus sp.	Sedge	0.16
Dichanthelium aciculare	Needle leaf witch grass	0.33
Dichaanthelium sp.	Witch grass	0.33
Diodia teres	Poor Joe	0.33
Eragrostis spectabilis	Purple lovegrass	0.167
Eupatorium compositifolium	Yankeeweed	0.67
Gaylussacia dumosa	Dwarf huckleberry	0.33
Haplopappus divericatus	Scratch Daisy	0.33
Hypericum gentianoides	Orange weed	0.33
Liatris tenuifolia	Shortleaf gayfeather	0.67
Licania michauxii	Gopher apple	0.33
Pinus clausa	Sand pine	0.16
Pteridium aquilinum	Brachen	0.15
Quercus laevis	Turkey oak	2.2
Schizachyrium sp	Little blue stem	1.5
Solidago tortifolia	Twisted leaf goldenrod	0.5
Stylisma patens	Coastal plain dawnflower	0.16
Vaccinium corymbosum	High bush blueberry	1
	Bare ground	66

Figure 15. Transect 2: Species Cover and Occurrence (Sand Pine Plantation)




Table 5. Transect 4. Species cover and occurrence (Sand Pine Plantation)

Date 11/5/09

Time: 10:30 Pm

Collector: David Clayton

Condition, Fair and cool

Wildlife observed: None

Fuel load: low


Community description: Sand pine plantation removed

East side Dykes Mill Pond and Green Head

<u>Scientific Name</u>	<u>Species</u>	<u>Percent Cover</u>
Aeschynomene americana	Aeschynomene	0.167
Andropogon virginicus	Broom sedge	0.33
Aristida stricta	Wire grass	12.3
Bulbostylis ciliatifolia	Capillary hair sedge	0.167
Chrysoma pauciflosculosa	Woody goldenrod	2.3
Chrysopsis lanuginosa	Lynn Haven goldenaster	1.5
Conyza canadensis	Canadian horseweed	1.6
Cyperus sp.	Sedge	0.83
Dichaanthelium sp.	Witch grass	1
Diodia teres	Poor Joe	2
Diospyros virginiana	Persimon	0.5
Eremochloa ophiuroides	Centipede grass	4
Eupatorium capillifolium	Dog fennel	13

Eupatorium mohrii	Mohr's thorough wort	1.8
Galactia sp.	Milk pea	0.167
Hypericum gentianoides	Orange weed	1.6
Ilex vomitoria	Yaupon	0.5
Pinus paulustris	Long leaf pine	0.83
Opuntia humifusa	Prickly pair cactus	1.67
Quercus hemaespherica	Diamond Oak	0.33
Rubus cuneifolius	Sand black berry	4
Schizachyrium sp	Little blue stem	1.3
Vaccinium arboreum	Sparkle berry	0.16
	Bare ground	48

Figure 16. Transect 4: Species Cover and Occurrence (Sand Pine Plantation)



UMAM Polygon I, Management Unit 12- Sand Hill

UMAM Polygon I, Management Unit 12, consists of 263.52 acres. This polygon is dominated by a sand hill community with an overstory dominated by turkey and live oaks with scattered remnant longleaf pine and an understory dominated by wire grass and a wide variety of herbaceous species. Reclamation activities within this upland community include re-introduction of fire, thinning of oaks to less than 150 trees per acre and planting of long leaf pine seedlings at a density not to exceed 200 trees per acre at final release. Fire was re-introduced to this area during the winter of 2004. A winter burn scheduled for the areas that had oak reduction. Prior to the re-introduction of fire, the dominant understory species was woody goldenrod. Oaks were thinned for the majority of Management Unit 12 in August of 2005. However, the portion of Management Unit 12 which contains Transect 5 was thinned in September of 2006. The re-introduction of fire and thinning of the turkey and live oaks have led to significant changes in the species composition. Two transects (transect #3 and #5) were located within UMAM Polygon I, Management Unit 12, and reflect baseline conditions (Table 6, 7 and Figure 16, 17).

In 2006, a total of 23 species were observed in transect 3 and 31 species in transect 5. A diverse understory of plants typical of sand hill vegetation was observed within each transect. No nuisance or exotic species cover occurred within these transects. The greatest cover class for each transect was bare ground with 47.5% (transect 3) and 68.5% for transect 5. Wire grass was the dominant vegetative species for both transects with 27.2 % cover for transect 3 and 22.2% cover for transect 5. A total of 12 species, Elliot's bluestem, wiregrass, Coastalplain honeycombhead, woody goldenrod, silver croton, witch grass, persimmon, pineland spurge, milk pea, pinweed, gopher apple and bracken fern were common to both transects.

In 2008, a total of 18 species were observed in transect 3 and 27 species in transect 5 slightly higher for transect 3 and lower for transect 5 than last year. A diverse understory of sand hill vegetation was observed again this year and no nuisance or exotic species were observed (Table 6, 7, Figure 16 and 17). The greatest cover class again was bareground with 36.1% cover for transect 3 and 37% for transect 5. The amount of bareground for each transect was greatly reduced and may be due to the re-introduction of fire. Wire grass was again the dominant vegetative species for both transects with 34% for transect 3 and 38% cover for transect 5. Wire grass cover increased by 6.8% for transect 3 and 15.8% cover for transect 5. A total of 9 species were common to both transects.

In 2009, a total of 15 and 29 species were observed within transects 3 and 5 respectively. The number of species is lower for transect 3 by three species and slightly higher by two species for transect 5. A diverse understory of sand hill vegetation was again observed during this monitoring event. No nuisance or exotic species were observed (Table 6, 7, Figure 17 and 18). The greatest cover class was bare ground for transect 3 with 44% cover while wire grass cover was the great percent cover with 42.5% cover for transect 5. Wire grass was again the dominant vegetative species for both transects with 39% for transect 3 and 42.8% cover for transect 5. Wire grass cover increased by 5% for transect 3 and 4.8% cover for transect 5.

Longleaf pines were planted in portions of UMAM polygon I, Management Unit 12 in the winter of 2004. However, longleaf pines were only observed in Transect 3 in 2006. A belt transects 600' feet in length and 30' feet in width was co-located with the vegetation transect. The number, height and condition of each planted tree were recorded. A total of 36 trees were observed or an average of 871 trees per acre. However, the winter burn in 2006 was extremely intense and killed nearly all planted pines. A total of 2 seedling pines were observed in 2007 both close to the ground and in the grass stage. During the 2008 monitoring, two planted pine seedlings were again observed, both in the grass stage. These areas were planted with less than 300 trees per acre during

the winter of 2008. During the 2009 sampling event, a total of 38 long leaf pines were observed along transect 3 and 24 along transect 5. Seedlings were in the grass stage and appeared healthy..

Interim success Criteria:

The interim success criteria have been met for UMAM I polygon I. Fire was re-introduced to the site, turkey and live oaks were thinned to less than 150 trees per acre and long leaf pine have been planted. In June of 2009, due to an increase in oak sprouts from the felled trees in the sand hills, ULW, and herbicide selective for oaks. It is hoped that this will reduce oak cover to less than the 150 oaks per acre required by the permit. No nuisance or exotic species occurred were observed within transects, fire adapted species dominate the vegetative cover, while wood species cover has been greatly reduced. Wire grass cover continues to increase and sandhill species dominate the polygons.

Table 6. Transect 3. Species cover and occurrence (Sand Hill)

Transect 3

Date 11/4/09

Time: 1:00 Pm

Collector: David Clayton

Condition, Fair and cool

Wildlife observed: Crow

Fuel load: Moderate

Community description: Sandhill with oaks thinned

Greenhead branch

Scientific Name	Species	Percent Cover	# speices
<i>Andropogon arctatus</i>	Pinewoods bluestem	3.3	1
<i>Andropogon virginicus</i>	Broom sedge	0.167	2
<i>Aristida stricta</i>	Wire grass	39	3
<i>Chrysoma pauciflosculosa</i>	Woody goldenrod	1.7	4
<i>Diospyros virginiana</i>	Persimon	0.167	5
<i>Gaylussacia dumosa</i>	Dwarf huckleberry	1	6
<i>Persea borbonia</i>	Red bay	1	7
<i>Pinus paulustris</i>	Long leaf pine	0.33	8
<i>Polygonella gracilis</i>	Wire weed	0.167	9
<i>Opuntia humifusa</i>	Prickly pair cactus	0.167	10
<i>Quercus incana</i>	Blue jack oak	0.167	11
<i>Quercus laevis</i>	Turkey oak	3.5	12
<i>Serenoa repens</i>	Saw palmetto	5.3	13
<i>Vaccinium darrowii</i>	Darrow's blueberry	0.167	14
<i>Vaccinium arboreum</i>	Sparkle berry	0.167	15
	Bare ground	44	

Figure 17. Transect Three Percent Cover




Table 7. Transect 5 Species and Occurrence (Sand Hill)

Transect 5

Date 11/5/09

Time: 8:50 am

Collector: David Clayton

Condition, Fair and cool

Wildlife observed: Wren, chipping sparrow


Fuel load: moderate due to herbicide of oaks

Community description: Sandhill

Scientific Name	Species	Percent Cover	# speices
<i>Andropogon arctatus</i>	Pinewoods bluestem	1.3	1
<i>Andropogon virginicus</i>	Broom sedge	0.33	2
<i>Aristida stricta</i>	Wire grass	42.8	3
<i>Asimina angustifolia</i>	Slimleaf paw paw	0.167	4
<i>Aster pilosus</i>	Frost aster	0.167	5
<i>Balduina angustifolia</i>	Coastalplain honeycomb head	1.5	6
<i>Baptisia lanceolata</i>	Gopherweed	0.167	7
<i>Bulbostylis ciliatifolia</i>	Capillary hair sedge	0.167	8
<i>Chrysoma pauciflosculosa</i>	Woody goldenrod	2.67	9
<i>Crataegus michauxii</i>	Michaux's hawthorn	0.33	10
<i>Dichanthelium sp.</i>	Witch grass	0.167	11
<i>Diospyros virginiana</i>	Persimon	2	12
<i>Eriogonum tomentosum</i>	buckwheat	0.167	13

<i>Gelsemium sempervirens</i>	Yellow jessamine	0.167	14
<i>Hypericum gentianoides</i>	Orange weed	0.5	15
<i>Ilex vomitoria</i>	Yaupon	0.167	16
<i>Licania michauxii</i>	Gopher apple	1.167	17
<i>Penstemon multiflorus</i>	Manyflowered beardtongue	0.167	18
<i>Pityopsis graminifolia</i>	Narrow leaf silkgrass	0.167	19
<i>Polygonella gracilis</i>	Wire weed	5.8	20
<i>Pteridium aquilinum</i>	Brachen	1.5	21
<i>Opuntia humifusa</i>	Prickly pair cactus	0.66	22
<i>Quercus incana</i>	Blue jack oak	0.167	23
<i>Scleria</i> sp.	Nutrush	0.167	24
<i>Smilax</i> sp.	Smilax	0.83	25
<i>Vaccinium corymbosum</i>	High bush blueberry	0.83	26
<i>Vaccinium myrsinites</i>	Shiny blueberry	1.3	27
<i>Yucca filamentosa</i>	Adam's needle	0.33	28
	Bare ground	33.5	

Figure 18. Transect 5: Species Cover and Occurrence (Sand Hill)



UMAM Polygon(s): VII, Management Unit 3- Planted Slash Pine Plantation

UMAM Polygon VII, Management Unit 3, consists of 11.5 acres of bedded planted slash pine that will be restored to a hydric pine flatwood. The overstory was dominated by planted slash pine. The shrub and understory was largely been shaded out by the near complete canopy closure of the slash pine. Pines were thinned to 225 trees per acre in 2007. Following the initial burn in the summer of 2005, it was determined that the shrubs could be kept to coppice sprouts with successive warm season burns. In winter 2011, wire grass tublings will be planted on 3' centers throughout the polygon.

In 2006, a total of 17 species were observed. The majority of the species were common to wet flatwoods. No nuisance or exotic species were observed. The greatest cover class observed was bare ground at 80.5%. The dominant vegetation was black ti ti with 6.5 percent coverage. The total shrub coverage was approximately 12%. No wire grass was observed within this polygon.

In 2007, a total of 18 species were observed, similar to baseline observations. The majority of the species were common to wet flatwoods. No nuisance or exotic species cover was observed. The greatest cover class was again bare ground with 77.3 percent cover. The slight increase in vegetative cover may be due to increased light reaching the understory since the dense pine canopy has been thinned. Swamp dog hobble had the greatest percent vegetative, each with 5 percent. Black titi cover was reduced from 6.5 % to 3.7%. This represents a reduction in black titi cover from the baseline observations. Overall shrub coverage within this polygon slightly increased from 12% in 2006 to 13.4% in 2007 and herbaceous cover has increased from last year. Wildlife observations included a blue jay, towhee, and cardinal.

In 2008, a total of 30 species were observed. The majority of the species were common to wet flatwoods. No nuisance or exotic species were observed. Bare ground again had the largest cover class with 58% down from 77.3% the year before. Black titi had the greatest cover class of the vegetation with 5.4%, increasing by 1.7%. Overall shrub cover within the polygon has increased from 13.4% in 2007 to 17% in 2008. Herbaceous cover also continues to increase over time. In 2008, herbaceous cover within the transect increased to 23.7%.

In 2009, a total of 28 species were observed. The majority of the species were common to wet flatwoods. No nuisance or exotic species were observed. Bare ground again had the largest cover class with 52% down from 58% the year before. Muscatine grape had the greatest cover class of the vegetation with 9.5%. Shrub cover within the polygon has decreased significantly from 17% in 2008 to 8.5% in 2009. Herbaceous cover also continues to increase over time. In 2008, the herbaceous cover increased to 30% and increase of 6.3% from the previous year.

Interim success Criteria:

Many of the management activities that will be used to restore UMAM VII, Management Unit 3 have been implemented and interim management activities completed or initiated. The forested canopy has been reduced to approximately 200 trees per acre (225) and is expected to further decline with the continuation of warm season burns. No exotic or nuisance species were observed. Herbaceous species cover is increasing and species present are consistent with wet pine flatwoods. A warm season burn was introduced in 2006, and the slash pines were reduced in density. The area will be planted with wire grass during the winter of 2009/2010 and a winter burn is planned for the winter of 2009/2010.

Transect 8 Hydric Pine flatwoods

Date 11/2/09

Time: 11:30 am

Collector: David Clayton

Condition, Fair and cool


Wildlife observed: Red shouldered hawk, titmouse, red bellied woodpecker

Community description: Wet flatwoods, slash pine thinned

Fuel load: low

Scientific Name	Species	Percent Cover	# speices
<i>Andropogon glomeratus</i>	Bushy bluestem	0.167	1
<i>Carex verrucosa</i>	Warty Sedge	0.5	2
<i>Centella asiatica</i>	Centella	0.8	3
<i>Chamaecrista nictitans</i>	Partridge berry	1.1	4
<i>Cliftonia monophylla</i>	Black titi	5.1	5
<i>Cyrilla racemiflora</i>	Red titi	0.167	6
<i>Dichantheium sp.</i>	Witch grass	3.16	7
<i>Diodia teres</i>	Poor joe	2.3	8
<i>Eupatorium capillifolium</i>	Dog fennel	1.5	9
<i>Gelsemium sempervirens</i>	Jessamine	0.33	10
<i>Hyper sp.</i>	St. John's wort	1	11
<i>Ilex glabra</i>	Gall berry	0.5	12
<i>Ilex vomitoria</i>	Yaupon	1	13
<i>Lachnanthes caroliniana</i>	Red root	6.5	14
<i>Leucothoe racemosa</i>	Swamp dog hobble	0.167	15
<i>Ludwigia decurrens</i>	Seedbox	0.5	16
<i>Lycopus virginicus</i>	Water horehound	0.3	17
<i>Myrica cerifera</i>	Wax myrtle	1.8	18
<i>Panicum sp.</i>	Panic grass	8	19
<i>Persea palustris</i>	Swamp bay	0.8	20
<i>Pinus palustris</i>	Longleaf pine	0.1	21
<i>Polygala lutea</i>	Candy weed	0.16	22
<i>Rhexia alifanus</i>	Savannah meadow beauty	0.06	23
<i>Rhynchospora microcephala</i>	Bunched beaksedge	0.167	24
<i>Rubus cuneifolius</i>	Sand black berry	1.3	25
<i>Smilax laurifolia</i>	Cat briar	0.167	26
<i>Vaccinium corymbosum</i>	Highbush blueberry	0.3	27
<i>Vitis rotundifolia</i>	Muscadine grape	9.5	28
	Bare ground	52	

Figure 19. Transect 8, Planted slash pine species cover and occurrence



(UMAM Polygon V, Management Unit 2, Hydric Pine Flatwoods

UMAM Polygon V, Management Unit 2 consists of 165 acres of fire suppressed shrub dominated hydric pine that will be restored to a hydric pine flatwood. The overstory is dominated by a near impenetrable shrub layer with a largely lacking tree canopy and herbaceous layer. Reclamation activities within this polygon include removal of shrub overstory with a Gyro-trac followed by continued treatment with selective herbicides if necessary, re-introduction of fire, planting of longleaf and slash pine trees at a rate of 436 trees per acre, planting wiregrass tubelings on 3' centers, and monitoring for nuisance / exotic plant species. If the seed bank does not respond, additional keystone flatwood species will be introduced as tublings.

Fire was re-introduced into this polygon during the summer of 2005. Two transects, 6 and 7 were established in different portions of the hydric pine flatwoods. The warm season burn was effective in reducing the overstory of shrubs in transect 7, however, by the time of the initial sampling event, the majority of the shrubs had sprouted from the roots and already formed an extremely dense shrub layer approximately 3-4' in height. The fire was less effective in the area surrounding transect 6. Many of the black ti ti within this transect did not burn.

In 2006, a total of 14 species were observed within the transect 6 and 16 in transect 7. Seven species were common to both sites, and all were shrubs. Both sites were dominated by shrubs with little overstory and little to no understory species due to the extremely thick shrub layer. No exotic species were observed. The greatest cover class observed for both transects was black ti ti with 69.87 % cover in transect 6 and 31.77 percent cover in transect 7. No wire grass was observed within this polygon. One other shrub species Fetterbush (15.3%) had significant cover within transect 6, myrtle leaved holly (15.4%) had significant cover in transect 7. Little bare ground was observed in transect 6 (3.7%) while 11.5% bare ground was observed in transect 7.

In 2007, a total of 12 species were observed within transect 6 and 9 in transect 7. Transect 6 had a similar species composition to the baseline while transect 7 had significantly fewer species observed probably due to the gyrotrack. Seven species were common to both sites, and all were shrubs. Both sites were dominated by 3-3.5' shrubs though each had an herbaceous component. While this did not represent significant cover in transect 6, 3.7% cover in transect 7 was red root, and early colonizing wetland species. The greatest cover class for both transects was bare ground with 40.8% for transect 6 and 48.2% cover for transect 7. This represents a significant shift in cover from black titi to bare ground due to the gyrotrack. Black titi cover was also greatly reduced from nearly 70% to 14% in transect 6 and from 31.77% to 28.1% cover in transect 7. The relative minor decrease in black titi cover in transect 7 may be the result of the intense warm season fire in 2006. Fetterbush was the dominant species by cover in transect 6 while black titi remained the dominant plant species by cover in transect 7. Continued management activities will further reduce shrub coverage.

In 2008, a total of 24 species were observed within transect 6 and 16 in transect 7. This represents a 50% increase in transect 6 and 56% increase in species in transect 7. The seed bank along both transects has started to respond and herbaceous species not identified previously have emerged. A total of 10 new herbaceous species were observed along transect 6 and 7 new herbaceous species in transect 7. Shrub cover along transect 6 increased from 3.7% to 47.57% an increase of 43.87% and along transect 7 remained approximately the same 48.1% in 2007 to 48.38% in 2008. Shrub levels at each site were beyond acceptable levels. Test plots using selective herbicides that eradicate target shrubs without impacting the native understory showed great promise. In the test plots, shrub levels were reduced from near 50% cover to less than 15% with two applications. In 2009, these treatments will be expanded across the landscape.

In 2009, a total of 11 species were observed in transect 6 and 14 in transect 7. This is a significant drop in species observed along transect 6 but a similar number of species observed along transect 7. The drop in species along transect 6 may be due to a reduction in shrub species found in transect 6. Six shrub species found in transect 6 in 2008 were absent in 2009, probably from the selective herbicide treatment. However the number of shrub species remained constant along transect 7. Herbaceous species cover along transect 6 was slightly lower in 2009 with 7.1%, down from 10.4 %. Similarly along transect 7 herbaceous cover was down from 12.07% in 2008 to 10.45 in 2009. However wire grass cover increased from 2% for transect 6 and 7 in 2008 to 3.8% for transect 6 and 2.8% in transect 7 in 2009. Nuisance shrub cover along transect 6 decreased dramatically from 47.57% in 2008 to 23% in 2009. Similarly, shrub cover along transect 7 decreased from 43.87% cover in 2008 to 25.2 along transect 7. This represents about a 48 and 60% reduction in shrub cover over the last year respectively for transect 6 and 7.

Interim Success Criteria:

Most of the management activities were completed by 2007 for of the UMAM V, Management Unit 2. Fire was introduced in 2005 and a second site prep burn occurred in December of 2007. A gyrotrack was employed (April-July) to reduce the shrub cover to basal sprouts. Wire grass tublings and long leaf pine seedlings were planted in late December/January 2008. No exotic vegetation has been observed at anytime in this polygon. In 2008, the herbaceous species observed within the polygon greatly increased by more

than 50%, indicating that the seedbank was responding to the shrub reduction. Planted wire grass had about a 65% survival. During the last year wire grass survival dropped to about 45%. As available additional wire grass will be added to these areas. The polygon will was treated with selective herbicides in 2009 to target nuisance shrubs, followed by a cool season burn during the winter of 2009/2010. A total of 5 acres of toothache grass will be planted near Dry Pond and 38 acres of wet prairie wire grass will be planted adjacent to Dry Pond. In addition, 5 acres of mixed wet flatwood species will be planted adjacent to Dry pond during the winter 2009/2010. Overall, shrub cover has been significantly reduced. In addition, a masters student will be evaluating the seed bank within the wet flatwood areas over the next year to determine viability of the seed bank. In addition in future the District will be adding diverse wet flatwoods species at the time of planting the wire grass to ensure a diverse habitat emerges.

Table 9. Transect 6 Species Cover and Occurrence (Hydric Pine Flatwoods)

Date 11/3/09
 Collector: David Clayton
 Wildlife observed: titmouse
 Community description:Wet flatwoods

Time: 10:00 am
 Condition, Fair and cool
 Fuel load:low

Scientific Name	Species	Percent Cover	# species
<i>Aristida stricta</i>	Wire grass	3.8	1
<i>Cliftonia monophylla</i>	Black titi	4.6	2
<i>Cyrilla racemiflora</i>	Red titi	1	3
<i>Ilex glabra</i>	Gall berry	1.3	4
<i>Ilex myrtifolia</i>	Myrtle-leaved holly	2.6	5
<i>Lachnanthes caroliniana</i>	Red root	0.16	6
<i>Lyonia lucida</i>	Fetter bush	16.1	7
<i>Osmanthus americanus</i>	Wild olive	0.16	8
<i>Persea palustris</i>	Swamp bay	0.66	9
<i>Rhynchospora microcephala</i>	Bunched beaksedge	3.2	10
<i>Vaccinium corymbosum</i>	Highbush blueberry	0.167	11
	Bare ground	66.5	

Figure 20. Transect 6: Species Cover and Occurrence (Hydric Pine Flatwoods)





Table 10. Transect 7. Species and Occurrence (Hydric Pine Flatwoods)

Date 11/2/09
 Collector: David Clayton
 Wildlife observed: none
 Community description: Wet flatwoods
 Time: 10:00 am
 Condition, Fair and cool
 Fuel load: low

Scientific Name	Species	Percent Cover	# speices
<i>Andropogon glomeratus</i>	Bushy bluestem	0.67	1
<i>Aristida stricta</i>	Wire grass	2.8	2
<i>Cliftonia monophylla</i>	Black titi	18.3	3
<i>Cyrilla racemiflora</i>	Red titi	5.1	4
<i>Eupatorium capillifolium</i>	Dog fennel	0.5	5
<i>Gaylussacia dumosa</i>	Dwarf blueberry	0.8	6
<i>Ilex glabra</i>	Gall berry	0.5	7
<i>Ilex myrtifolia</i>	Myrtle-leaved holly	0.16	8
<i>Lachnanthes caroliniana</i>	Red root	5.5	9
<i>Leucothoe racemosa</i>	Swamp dog hobble	0.167	10
<i>Lyonia lucida</i>	Fetter bush	1.7	11

<i>Persea palustris</i>	Swamp bay	0.167	12
<i>Rhynchospora microcephala</i>	Bunched beaksedge	0.167	13
<i>Vaccinium corymbosum</i>	Highbush blueberry	1.5	14
	Bare ground	61.1	

Figure 21. Transect 7. Species and Occurrence (Hydric Pine Flatwoods)



UMAM Polygon V1, Management Unit 5, Inland Ponds and Sloughs

UMAM Polygon V1, Management Unit 5 consists of 24.880 acres of a dammed slough (Dykes Mill Pond) that will be restored to slough/marsh. The overstory for most of the area is absent though a fringe of cypress remains along the ponds edge. The majority of the area is dominated by water lilies and other aquatic submerged vegetation. Reclamation activities within this polygon include the removal of Dykes Mill Pond dam, and spanning the gap with railcar bridge, planting of cypress and black gum saplings and planting the area with herbaceous and shrub species, if after 2 years, the native wetland understory is < 50%. Dykes Mill Pond was removed in August of 2006 and bridge construction completed in April 2007. With the removal of the dam there have been great changes to the pond. By September 2007 most of the pond had evaporated leaving only small flooded areas. Wet prairie vegetation has greatly spread across the newly exposed sediments and a braided stream channel has emerged across most of the previously flooded area. Sampling last year occurred from a canoe while this year I was able to walk across the entire pond.

In 2006, a total of 7 species were observed within transect 9. The species were common to freshwater marshes within the region. No exotic species were observed. The dominant species observed was fragrant water lily with 45 % cover. Florida yellow bladderwort was also common with 19.2 % cover. Open water

was common with 34% cover, indicating that much of the transect occurs in what is currently a pond. Wildlife was observed included wood ducks and a great egret.

In 2007, a total of 11 species were observed within transect 9. Species were common to wet prairies with some minor freshwater marsh species. This represents a major shift in species composition and reflects the shift from an aquatic to wet prairie. No exotic species were observed. Fragrant water lily cover was greatly reduced from 45% in 2006 to 3.23% cover in 2007. Florida yellow bladderwort was not observed within the transect and open water was also greatly reduced from 34% cover to 2.2 % cover. Another significant occurrence was the cover of bare ground which did not exist in 2006, but represented 41% of the cover in 2007. The two dominant plant species were horned beaksedge with 30% cover and a beaksedge that was not in flower with 12% cover, both species common to wet soils and not tolerant of aquatic systems. A species of note, *Drosera intermedia* (Water Sundew) a state threatened species was commonly observed. Wildlife observations included a pair of sandhill cranes (State Threatened species), fresh hog tracks, little blue heron, great egret, and chipping sparrows.

In 2008, a total of nine species were observed along transect 9. Followed by two years of drought, Dykes Mill ponds water level came up flooding most of the historic foot print. The removal of the dam reduced the water level by approximately 6' but not the expected 20 to 30'. The transect was flooded with 6" to 3' of water. Water lilies and aquatic vegetation abound and are thriving providing important habitat for wildlife. An alligator nest was observed along the bank and baby alligators were observed with their 6' mother during sampling. Open water was the dominant cover class with 46.3 percent cover. The dominant vegetative species was fragrant water lily with 33 percent cover.

In 2009, a total of 9 species were again observed along transect 9. Water levels remained high flooding most the historic footprint. Since reclamation activities took place Dykes Mill Pond water levels appear to be more or less stable approximately 6' lower along the shore line than were observed during the baseline condition. The entire transect length was inundated with 4" to 3' of water. Water lilies and aquatic vegetation abound and are thriving providing important habitat for wildlife. Open water was the dominant cover class with 38 percent cover. The dominant vegetative species was fragrant water lily with 35.3 percent cover

Interim Success Criteria:

Most of the management activities used to restore UMAM VI, Management Unit 5 have been completed. The archeological study was completed and the dam removed in August of 2006. The new bridge was completed in April of 2007. Since the removal of the dam the pond drained during the drought and much of the dry pond area was dominated by grasses and sedges. In 2008, the water levels increased due to the end of the drought and a shallow pond formed in 2008. Cypress trees and black gums were planted along the edges of this system in the spring of 2007/2008. The shrub areas adjacent to Dykes Mill Pond were Gyro-Trac'd in 2007. Due to increasing shrub cover in the area, selective herbicides will be used to reduce shrub cover.


Table 11. Transect 9. Species and Occurrence (Slough / Marsh)

Date 11/6/09 Time:9:30 am
 Collector: David Clayton Condition, Fair and cool
 Wildlife observed:Great white egret, little blue heron, alligator
 Community description: Marsh Fuel load: N/A

Scientific Name	Common Name	Percent Cover	# species
-----------------	-------------	---------------	-----------

Drosera intermedia	Water sundew (Fl threatened)	0.16
Eleocharis cellulosa	Eleocharis	3.3
Lachnanthes caroliniana	Red root	0.5
Nuphar advena subsp. Orbiculata	Spatterdock	4.7
Nymphaea odorata	Fragrant water lily	35.5
Rhynchospora inundata	Horned beaksedge	6.8
Utricularia floridana	Florida bladderwort	1.2
Websteria confervoides	Websteria	7.5
Xyris sp	Yellow-eyed grass	2.3
	Open water	38

Figure 22. Transect 9. Species and Occurrence



Qualitative Monitoring

Materials and Methods

Qualitative vegetation monitoring will include assessment of the vegetation, both ground cover and planted trees, wildlife use observations, and general habitat health. Pedestrian surveys increase site coverage and include a 30+ minute meandering walk-path intended to provide information useful in management and to

determination the success of management activities. A walk path traversed as much habitat as possible. The pedestrian walk-path continued as long as species were being added, however, once additional species were not recorded for 3 minutes the survey was complete. Representative photos and a community description and health were provided for each walk-path. Fuel load for each habitat was determined and the presence of any threatened or endangered species were recorded. Plants were listed in the data sheet in the following categories (tree, shrub, vine or herbaceous) to give a better understanding of composition of the habitat. Wildlife observations were also recorded for each walk-path (Figure 13) provides the location and coverage of transects and the data sheets can be found in (Appendix 4).

Results and Discussion

A total of 13 pedestrian transects were located at the SHLMB (Figure 13) Three pedestrian surveys were located in Management Unit 1, portions of UMAM Polygon IV, one in Management Unit 2, UMAM Polygon V, one in Management Unit 4, portions of UMAM Polygon IV, four in Management Unit 10, Polygon III, three in Management Unit 12, UMAM Polygon I, and one in Management Unit 14, portions of UMAM Polygon IV (Appendix 7).

Management Unit 1, UMAM Polygon IV, Preserved High Quality Forested and Herbaceous Wetlands

Management Unit 1, UMAM Polygon IV consists of 574.839 acres of a wide variety of preserved wetland habitats including approximately FLUCCS: 621 – Cypress, 617 – Mixed Wetland Hardwoods, 644 – Emergent Aquatic Wetlands, 611 – Bay Swamps, 641 – Freshwater Marshes, 616 – Inland Ponds and Sloughs, 640 – Vegetated Non-Forested Wetlands and 643 – Wet Prairies. The management goal for this polygon is the preservation of the existing high quality wetlands. Two of the pedestrian survey paths (M8 and M9) in Management Unit I, UMAM Polygon IV, were located in cypress dominated wetlands, while the third pedestrian survey path (M10) was located in an overgrown hydric pine flatwoods. However it is suggested that this transect be kept but the designation and analysis changed to the more appropriate Management Unit 2, UMAM Polygon V.

In 2006, a total of 38 species were observed in M8, while 32 species were observed in M9. Twenty nine of the species were common to both transects. Five tree species were observed in M8 while 3 tree species were observed for M9. Eight and nine shrub species were observed in M9 and M8 respectively, though cover of shrubs was not significant. Twenty one herbaceous species were observed in M8, while 19 herbaceous species were observed in M9. No nuisance or exotic species were found in M8, though a small patch of torpedo grass was observed in M9. Fuel load was low for each area and no threatened or endangered species were observed. Water levels in both areas were extremely low due to the drought and many of the herbaceous species such as pickerel weed, duck potato (*Sagittaria latifolia*) and fragrant water lily (*Nymphaea odorata*) had browned or appeared dead. Cypress seedlings were numerous in both areas. Wildlife was abundant.

In 2007, a total of 39 species were observed for M8 similar in number to last year four new species, bushy bluestem, beauty berry, sweet pepperbush, and pale meadow beauty were observed. These were observed in the normal pool area and germinated due to the prolonged drought that has left the lake beds dry. Three species previously observed, water shield, bog buttons, and bladder wort were not observed, primarily due to the absence of an aquatic habitat. Along M9, a total of 31 species were observed, again similar in number to last year. However 8 species were not observed this year and include water shield, clustered sedge, Virginia willow, silver bay, pickerel weed, duck potato, bladderwort and yellow eyed grass. These are primarily aquatic species and were not found on the dry lake beds. Nine additional species were observed including bushy

bluestem, sedge, black titi, witch grass, yaupon, sweet gum, savannah meadow beauty and American cupscale. The new species with the exception of the American cupscale are facultative wet species that have invaded the dry lake beds. Shrub cover for both transects was very low. No nuisance or exotic species were observed. Fuel load was low for each area and no threatened or endangered species were observed. Water levels in both areas were extremely low due to the drought and many of the herbaceous species aquatic species were absent. A wildfire occurred within this polygon and destroyed approximately 12 acres of cypress by burning the roots and occasionally the trunk of the cypress. Details on the wildfire have been recorded in the Fire Management section. Aside from the continued drought this polygon is very similar to last year.

In 2008, a total of 42 species were observed in M8 and increase of 3 species, swamp dog hobble, swamp laurel oak and savannah meadow beauty (Appendix 4).

. Bladderwort and bog buttons were again not observed in these areas along with water shield and Marsh St. Johns Wort. The area is starting to recover from the prolonged drought. Water levels were about ½” above the soil surface and aquatic plants were starting to emerge. Along transect M9, a total of 36 species were observed, a slight increase from the previous year (Appendix 4). Water levels were starting to increase in this area with the average water depth approximately 2” in depth. The species observed were transitional or upland species that had germinated in the wetland during the drought and included groundsel tree, winged sumac and pine barrens goldenrod. As the system recovers and water levels increase it is expected that the system will recover and the upland species will be removed by increasing water levels. No nuisance or exotic species were observed. Fuel load was low for each area and no threatened or endangered species were observed. Water levels in both areas were extremely low due to the drought and many of the herbaceous species aquatic species were absent. Wildlife observed included chipping sparrows, southern cricket frogs, a kingfisher and a red bellied wood pecker.

In 2009, a total of 39 species were observed along M8, probably a result of recovering from the drought. Water shield and St. John’s wort were again observed now that the water has returned. Dog fennel, red root, and centella were again observed within the area. Water levels were between 12” and 14”. Along M9, a total of 44 species were observed a marked increase from the low of 36 from the previous year. Many of the wetland herbaceous species absent in the drought have returned and are flourishing once again. No nuisance or exotic species were observed in this area nor any threatened or endangered species.

Interim Success Criteria:

Interim success criteria have been met and include exotic vegetation cover < 2% per acre, nuisance vegetation cover < 5% per acre, and maintaining or improving in ecological function. Water levels have recovered and both areas had an increase in wetland dependant species.

Management Unit 2, UMAM Polygon V, Hydric Pine Flatwoods

Management Unit 2, UMAM Polygon V consists of 146.678 acres of FLUCCS 635 hydric pine flatwoods. The management goal for this polygon includes the enhancement and restoration of the degraded hydric pine flatwoods. Two pedestrian transects (M10 and M11) were located in Management Unit 2, UMAM Polygon V. Both of these areas are overgrown, degraded hydric pine flatwoods dominated by a variety of tree and shrub species. Both areas were burned during the summer of 2005, though fuel loads in both areas are moderate and additional fires are warranted. Dominant species cover along M10 was black ti ti with some silver bay and slash pine, while M11 was moved slightly in 2007 to better reflect the wet flatwoods. The previous transect was located in a mixed bayhead. Wire grass was present in M11, but absent in M10.

In 2006, a total of 32 species (8 trees, 17 shrubs, 4 vines and 3 herbaceous species) were observed along M10.

In 2007, shrub reduction was completed in both areas using a gyrotrack. Shrubs were thinned in June and the areas were burned in December 2007. A total of 40 species were observed in along M10 while 16 species were observed in M11. No nuisance exotic species were observed in either area. The increase in species along M10 may be due to increased access to the area due to the gyrotrack and the fact that the site is more of a mixture of wet flatwoods with species from an adjacent bayhead. Successive fires should remove the bayhead species. A total of 22 species were observed along M11 in 2007. The lower number of species found in M11 is more reflective of a site that had been overgrown with shrubs and recently reduced to ground level by the gyrotrack. Over time it is expected a greater number of species will germinate from the seed bank. Wildlife observed included robin, kingfisher, black vulture, phoebe, anole and cardinal.

In 2008, a total of 51 species were observed, eleven species more than the year before. The seed bank has started to respond in this area and additional species observed were primarily herbaceous species commonly found in wet flatwoods. Shrubs in this area had also increased in cover and will be targeted in the coming year with selective herbicides to reduce shrub cover while preserving the understory vegetation. A total of 26 species were observed along M11 in 2008, an increase of 4 species. No nuisance or exotic species were observed during the 2008 sampling. A minor amount of hog damage was observed adjacent to polygon, and trappers have been notified. Wildlife observed included titmouse, red bellied wood pecker, flicker, blue jay and raccoon tracks.

In 2009, a total of 54 species were observed within the meandering transect of M 10. This represents an increase of 3 species. Several shrub species that had been a problem in the area were not observed in this year's sampling. The species observed were common to wet flatwoods. Targeted shrub densities have greatly decreased in these areas and wire grass will be planted in the winter of 2009/2010. A total of 28 species were observed along M11, a slight increase from the previous year. This area continues to develop, and additional herbicide work is needed here to insure that the shrub density continues to decline. Positioned adjacent to the Green Ponds, this area should have good natural recruitment. No nuisance or exotic species were observed. Wildlife observed included deer tracks and a titmouse.

Interim Success Criteria:

Interim success criteria include exotic vegetation cover < 2% per acre, nuisance native vegetation cover < 5% per acre, increasing herbaceous groundcover, decreasing density of woody shrub layer, planted pines are surviving and healthy and prescribed burns have been conducted in accordance with fire management plan. The interim success criteria have been met for this polygon. No nuisance exotic or nuisance native species cover has been observed, and the prescribed burns have been conducted in accordance with the fire management plan. Shrubs were reduced to ground level in both areas using a gyrotrack and both areas and herbaceous vegetation cover is increasing within the polygon. These sites were burned in December 2007 and will be burned again in 2009. Positioned adjacent to the Green Ponds, this area should have good natural recruitment. Wildlife observed included deer tracks and a titmouse. Due to the numbers of existing pine trees this area will not need supplemental tree planting, however, wire grass will be added to the polygon in 2009.

Management Unit 10, UMAM Polygon III, Xeric and Live Oak

Management Unit 10, UMAM Polygon III consists of 493.852 of FLUCCS 421 – Xeric Oak and 427 – Live Oak. Management goals include the preservation and the re-introduction of fire to upland sandhill communities dominated by oaks. Management activities include the introduction of fire using dormant season burns, and the eventual introduction of growing-season burns (anticipated 3 to 5-year and 5 to 7-year burn cycles), and the reduction of oak in portions of management unit as selected by QMS (Qualified Mitigation Supervisor), and monitoring for nuisance / exotic plant species. Other management activities may include the supplemental planting of longleaf pine (436 trees per acre) and wiregrass (6' centers or direct seeding as 2-5 pounds per acre

as determined by the QMS. Live and turkey oaks were selectively harvested from portions of Management Unit 10, UMAM Polygon III in September of 2006. As a result the fuel load is high for most of these areas and a prescribed burn is scheduled for a dormant season burn in the winter of 2008/2009. Good coverage of wire grass was observed throughout Management Unit 10 so no additional planting will be required. Initial burns for portions of Management Unit 10 were conducted during the growing season. Wire grass was observed in flower for these areas. Continued warm season burns should ensure an increasing cover of wire grass throughout the polygon. Four transects were located within Polygon 10, M1, M2, M12 and M13).

In 2006, two transects M1 with 44 species (9 trees, 5 shrubs, 3 vines and 27 herbs) and M13 with 54 species (9 trees, 6 shrubs, 2 vines and 37 herbs) were species rich, while M2 with 29 species (6 trees, 6 shrubs, 3 vines and 14 herbs) and M12 with 26 species (12 trees, 3 shrubs, 3 vines and 8 herbs) were generally lacking a diverse herbaceous cover. This may be due to the shading of the understory by overstory oaks. However, all of the transects had between 19 and 35 species in common. Scattered diamond oak and sand pine may also be reflective of a historic lack of fire. No nuisance exotic coverage was observed, though a small patch of Bahia grass was found at the gate adjacent to the road for the transect M1. In the transect M1, a Florida threatened species Gulf coast lupine (*Lupinus westianus*) was located throughout the sand hill upland while smooth barked St. John's wort, a Florida Endangered species, was located adjacent to the solution pond 1. Gopher tortoise burrows were observed along pedestrian transects M12 and M13.

In 2007, two transects, M1 was observed with 67 species (10 trees, 16 shrubs, 3 vines and 38 herbaceous species) while, along M13 62 species (9 trees, 7 shrubs, 3 vines and 43 herbaceous species) was observed (Appendix 4). Along M2 38 species (8 trees, 5 shrubs, 2 vines and 23 herbaceous species) were observed and 34 species were observed along transect M12 (13 trees, 4 shrubs, 3 vines and 14 herbaceous species) (Appendix 4). M1 had 5 newly observed species and 3 species were not observed in 2007 and were sky blue lupine, bladderwort and yellow eyed grass. Ten new species were observed along M13 and two species, dwarf huckleberry and bracken fern were not observed. Along M2 13 additional species were observed while, 4 species Florida jasmine, red chokeberry, pale meadow beauty and lopsided Indian grass were not observed. Finally, M12 also had 13 additional species observed while 5 species were not observed and included American holly, gopher apple, sand pine, shiny blueberry, and Adam's needle. The observation of additional species may be due to increased scrutiny of the polygon and habitat improvement due to successive fires. Aside from a small patch of Bahia grass at the entrance to M1 no nuisance or exotic species were observed. Gulf coast lupine was observed at two transects, M1 and M13. Sand pine and Florida jasmine may have been removed by earlier fires. The habitat all appears healthy and vigorous. These areas were burned during the winter burns in December of 2007. Wildlife observed included a downy woodpecker, pileated woodpecker, raccoon tracks, otter tracks, gopher tortoise, deer tracks, turkey tracks, cardinal, towhee, titmouse and mockingbird.

This polygon is represented by four transects, M1, M2, M12 and M13. In 2008, M1 was observed with 69 species, 2 species greater than in 2007. M2 was observed with 35 species, three fewer than the previous year. In 2008, M12 was observed with 44 species, 10 species greater than in 2007 and M13 was observed with 59 species, three species fewer than in 2007. Species observed were typical of sand hill species. Gulf coast lupine was again observed in this location and is thriving in M1, M2 and M13. Wire grass continues to thrive in these areas. No nuisance or exotic species were observed, except for a small area at the entrance to M1.

This polygon is represented by four transects, M1, M2, M12 and M13. In 2009, M1 was observed with 74 species, 4 species greater than in 2007. This area is recovering from the drought and many of the species not observed adjacent to the pond have been seen again with the filling of the pond. The associated uplands are increasing in fuel and will be burned in 2010. M2 was again observed with 35 species. This area is managed under a 5-7 year burn cycle and as fuels increase perhaps species may become less common that require more light. A total number of 48 species were associated with M12, four greater than the previous year. A selective herbicide was used in this area to reduce the cover of hardwoods and help release the wire grass. Several hard

wood species cover was dramatically reduced in this area. Along M 13, 64 species, five species greater than in 2008 were observed. Species observed were typical of sand hill species. Gulf coast lupine was again observed in this location and is thriving in M1, M2 and M13. Wire grass continues to thrive in these areas. No nuisance or exotic species were observed, except for a small area at the entrance to M1.

Interim Success Criteria:

Interim success criteria have been met, no nuisance native or exotic vegetation have been observed, except for a small patch and M1's entrance. Diverse is good and continued fire within these areas will ensure a diverse sand hill community. Wire grass cover is good to excellent and oaks have been thinned.

Management Unit 11, UMAM Polygon II, Upland Slash or Sand Pine Plantations

Management Unit 11, UMAM Polygon II consists of 383.484 acres of FLUCCS 411 Longleaf Pine / Wiregrass restored from slash or sand pine plantations. The restoration goal for this area is to restore the sites to a sand hill community from a sand or slash pine plantation. Management activities will include the re-introduction of growing season burns, removal of planted pines, re-planting with 436 long leaf pine seedlings per acre and if needed the addition of wire grass tublings or seeding. Initial fire was introduced to the slash pine areas in 2005, while site prep burns will take place in the winter of 2008 for the previous sand pine areas. Trees were harvested from April to November 2007. One transect (M5) was located within Management Unit 11, UMAM Polygon II. This area had already undergone a warm season burn that greatly reduced the shrub cover. Overstory was removed in April 2007. Much of the understory was in fairly good condition with good diversity typical of the sand hills.

In 2006, a total of 50 species (6 trees, 7 shrubs, 2 vines, and 35 herbaceous species) were observed. Wire grass was the dominant grass species within the area. However, the emerging shrub layer was dominated by diamond oak.

In 2007, a total of 49 species were observed (7 trees, 8 shrubs, 2 vines and 32 herbaceous species) (Appendix 4). Nine new species were observed while 10 species initially present were not observed. The changes in species composition may be due to the tree harvest which greatly disturbed the understory. Species were common to the sandhill community. Wire grass was common and appeared to be the dominant species. Much of the shrub layer was reduced to sprouts and much of the diamond and live oak was destroyed during the tree harvest. A site prep burn is planned for winter 2008. Wildlife observed included: cardinal, fence lizard, titmouse, mourning dove, fish crows, red bellied wood pecker, phoebe, squirrel, two deer and lined race runner.

In 2008, a total of 59 species were observed (7 trees, 8 shrubs, 3 vines and 41 herbaceous species). Three additional species were observed. These were all typical dry sand hill species. Changes in composition are likely due to increased light from harvest of slash pine, followed by the initial burn. Wire grass continues to thrive and flowered this year. The area is returning to a sand hill from the planted pine plantation. Diversity will probably continue to increase due to increased fire rotations and response of the seed bank. No wild life was observed during sampling.

In 2009, a total of 61 species were observed (7 trees, 8 shrubs, 3 vines and 43 herbaceous species). Three additional species were observed. These were all typical dry sand hill species. Wire grass continues to increase in cover. The area has increased litter and will be burned during the winter of 2009/2010. The area is returning to a sand hill from the planted pine plantation. Diversity will probably continue to increase due to increased fire rotations and response of the seed bank.

Interim Success Criteria:

The interim success criteria have been met within this transect. No nuisance or exotic species were observed. Wire grass is the dominant species. The ground cover is diverse and typical of a sandhill. Diversity will likely continue to increase as the seed bank responds and with frequent fires.

Management Unit 12, UMAM Polygon 1, Sand Hill

Management Unit 12, UMAM Polygon 1 consists of 263.52 acres of FLUCCS: 411 – Longleaf Pine / Wiregrass (Mesic Pine Flatwoods) restored from 421 –Xeric Oak habitat.

The goal for this polygon is to restore a diverse sand hill. Restoration activities include the re-introduction of growing season burns, removal of oak ≤ 12 inches DBH and herbicide treatment of stumps, planting of longleaf pine (436 trees per acre), and monitoring for nuisance / exotic plant species. Oak eradication was conducted for Management Unit 12, UMAM Polygon 1 during the summer of 2005. . Fire was re-introduced in 2004 to the polygon and cover of the once dominant shrub woody goldenrod has been greatly reduced. Since the initial fire, two additional fires, the most recent in December of 2007, further reduced shrub and woody goldenrod cover. Wire grass has flowered for two consecutive years in most of this habitat. Wire grass is again the dominant herbaceous species within this polygon. The sand hill habitat within this polygon is very diverse and considered high quality with an excellent herbaceous species composition. The majority of the polygon was planted with longleaf pines in 2004, however, several areas on the north side of Green Head Branch will be re-planted with in 2008. Two transects (M3 and M4) were located within this polygon.

In 2006, a total of 35 species (7 trees, 2 shrubs, 2 vines, and 24 herbs) were observed along pedestrian transect M3, while 68 species (8 trees, 9 shrubs, 2 vines and 49 herbs) were observed within M4. The species were typical of the sand hill though in wetter areas of M4 adjacent to Little Deep Edge Pond, more pine flatwood vegetation occurred. Small patches of centipede grass were observed along the pedestrian transect M3.

In 2007, a total of 42 species (8 trees, 7 shrubs, 2 vines and 25 herbs) were observed along M3 (Appendix 4). The additional, shrub species observed may be due to expanding the path further to the west and up an old ridgeline. Shrubs were typical of the sandhill and high in wildlife value. Twelve new species were observed within this transect and may again be due to expanding the pedestrian survey. Five species, Southern magnolia, golden aster, bracken fern, Carolina milkweed and pinewoods milkweed were not observed in this years sampling. This may be due to the later sampling when some of these species are less noticeable following flowering and fruiting. Along the pedestrian transect M4 a total of 69 species (8 trees, 11 shrubs, 2 vines and 48 herbaceous) species were observed. A total of 17 new species were seen this year and 13 species previously observed were not seen this year. The area surrounding this pedestrian meander was burned during the winter of 2007 and the fire was particularly hot killing some turkey and live oaks and also may have removed some of the less fire tolerant species. Centipede grass which was observed as a minor component in the polygon was completely absent following the fire. Another species apparently removed by the fire was the slender crab grass. Other new species may have emerged from the seedbank once the fire exposed bare ground. Wildlife observed within this polygon included rabbit and raccoon tracks, and an active gopher tortoise burrow. In addition several threatened and endangered species were observed including southern crab apple, smooth barked St. John's wort and Gulf Coast lupine.

In 2008, a total of 53 species were observed along M3, an increase of 11 species from the previous year. The additional species were common to sand hills and species number may be increasing as a result of the shrub layer reduction due to successive fires. M4 is the most diverse of the areas of the bank. A total of 87 species were observed within this area. This is an increase of 18 species and may be due to a recent prescribed fire. Oak and shrub densities are low in this area and there are pockets of wet flatwoods within the sand hill vegetation. As shrub levels are reduced and continued fires will help in keeping the observed diversity. No nuisance or exotic species were observed though small patches of centipede grass were observed by staff. Some

expansion of turkey and live oaks were observed within these areas and will be reduced if cover continues to expand. Wildlife observed within this area included turkey and raccoon tracks, active gopher tortoise burrow, down wood pecker, chickadee. Threatened and endangered species include gulf coast lupine, crab apple and smooth barked St. Johns wort adjacent to cat pond.

In 2009, a total of 55 species were observed along M3, an increase of 2 species from the previous year. This area is quite diverse and additional species were common to sand hills and species number may be increasing as a result of the shrub layer reduction due to successive fires. In order to reduce oak sprouts from felled trees, the area was treated with Velpar (ULW), an herbicide selective for hardwood species. The area will be burned in the winter of 2009/2010. It is expected that Velpar will reduced the hard wood cover to below 150 trees per acre and keep the sand hill open and park like. M4 is the most diverse of the areas of the bank. This area historically had a low density of pines and oaks and continued fires should keep this area in excellent condition. A total of 91 species were observed within this area, and increase of 4 species. This area has an abundance of late successional sand hill species and quality of habitat in this area is very high. Oak and shrub densities are low and there are pockets of wet flatwoods within the sand hill vegetation. As shrub levels are reduced and continued fires will help in keeping the observed diversity. No nuisance or exotic species were observed though small patches of centipede grass were observed by staff. Some expansion of turkey and live oaks were observed within these areas and will be reduced if cover continues to expand. Threatened and endangered species include gulf coast lupine, crab apple and smooth barked St. John's wort adjacent to cat pond.

Interim Success Criteria:

This polygon has reached the restoration goals set forth in the interim success criteria. The three controlled burns within this polygon have greatly reduced the cover of woody golden rod and stimulated the cover of wire grass and other grasses and forbs. Oaks have been reduced to less than 150 trees per acre and the herbaceous vegetation is dominated by wire grass. Longleaf pines have been planted throughout the polygon in winter 2005.

Management Unit 14, portions of UMAM Polygon IV, Lakes

Management Unit 14, portions of UMAM Polygon IV consists of 164.958 acres of FLUCCS 520, lakes. The goal for this polygon is the preservation of the lake and aquatic habitat. One pedestrian transect (M6) was placed within the polygon around Garret Pond. The water levels at Garret pond were very low due to the summer drought. Much of the lake bottom was exposed and had been colonized by a variety of wetland grasses and sedges. Two small pools remained.

In 2006, a total of 36 species (5 trees, 7 shrubs, 1 vine and 23 herbs) were observed. Vegetation was typical of a diverse pond within the region. A small patch of torpedo grass was observed at the boat ramp to the pond. A zone of Smooth barked St. John's wort and seedlings was observed just below the shrub layer surrounding the pond. Some species such as pickerel weed appear to have been set back by the drought and most of the leaves and stem have browned.

In 2007, a total of 24 species were observed (5 trees, 7 shrubs, 1 vine and 11 herbs) (Appendix 4). Due to the extended drought, this pond has been dry for approximately a year. The reduction in herbaceous species is due to the lack of water. Most of the absent species were aquatic or required wet conditions to thrive. The small patch of torpedo grass at the old boat launch had been sprayed during the summer and none was observed during the fall sampling. Dog fennel has continued to invade the site and many of the aquatic species were absent.

In 2008, a total of 20 species were observed (5 trees, 7 shrubs, 1 vine and 8 herbs) (Appendix 4). Due to the extended drought, this pond has been dry for approximately 2 years. The reduction in herbaceous species is due

to the lack of water. Most of the absent species were aquatic or required wet conditions to thrive. Dog fennel has continued to invade the site and several wet flatwood species have been observed. No torpedo grass was observed during the fall monitoring.

In 2009, a total of 35 species were observed (5 trees, 7 shrubs, 1 vine and 23 herbs). The drought has ended and Garret Pond is again filling with water. Many of the wetland dependant species are once again flourishing within the pond and shoreline. The dog fennel and invading upland species have been drown and are being replaced by wetland vegetation. No torpedo grass was observed during the fall monitoring.

Interim Success Criteria:

No nuisance or exotic species were observed. Since the end of the drought, the water has returned to Garret Pond and spatter dock has re-emerged from the sediments. The site appears to be maintaining normal ecological functions and wetland vegetation again thrives in the pond.

Qualitative Field Assessment Form

Date: 10/30/2009 Time: 10:00 am Data Collector: David Clayton
 Location: Pedestrian Transect # M1 near photo point 15
 Management Unit: 10

Nuisance Species: Bahia grass at gate entrance

Fuel Load: Moderate

Wildlife Observations: wren, towhee

T& E Species: Moderate population of Gulf Coast Lupine in sand hill and Smooth Barked St. John's Wort around pond

Community Description: Sandhill upland adjacent to a solution pond. Sandhill with good diversity and excellent groundcover. Marsh zonation still present, pond dry most of year, center full of dog fennel.

Scientific Name	Common Name	Tree	Shrub	Vine	Herb
<i>Agalinis setacea</i>	Threadleaf false foxgloves				X
<i>Amphicarpum muhlenbergianum</i>	Blue maidencane				X
<i>Andropogon glomeratus</i>	Busy blue stem				X
<i>Aristida stricta</i> var. <i>beyrichiana</i>	Wiregrass				X
<i>Asimina angustifolia</i>	Slim leaved paw-paw				X
<i>Baptisia lanceolata</i>	Gopher weed				X
<i>Baulduina angustifolia</i>	Coastal plain honeycombhead				X
<i>Bulbostylis ciliatifolia</i>	Cappillary hairsedge				X
<i>Ceanothus microphyllus</i>	Redroot				X
<i>Centella asiatica</i>	Centella				X
<i>Cephalanthus occidentalis</i>	Button bush		X		
<i>Chrysoma pauciflosculosa</i>	Woody Goldenrod				X
<i>Cliftonia monophylla</i>	Black ti ti		X		
<i>Crysopsis scabrella</i>	Goldenaster				X
<i>Cyrilla racemiflora</i>	Titi		X		
<i>Dalea pinatta</i>	Summer farewell				X
<i>Dicanthelium scoparium</i>	Panic grass				X
<i>Dicanthelium spp.</i>	Panic grass				X
<i>Diospyros virginiana</i>	Persimon	X			
<i>Eleocharis sp.</i>	Eleocharis				X
<i>Eriogonum tomentosum</i>	Wild Buckwheat				X
<i>Eupatorium capillifolium</i>	Dog fennel				X
<i>Eupatorium compositifolium</i>	Yankee weed				X
<i>Eupatorium mohrii</i>	Eupatorium				X
<i>Euphorbia inundata</i>	Florida pineland spurge				X
<i>Euthamia caroliniana</i>	Flat-topped goldenrod				X
<i>Galactia sp.</i>	Milk pea				X
<i>Gaylussacia dumosa</i>	Dwarf huckleberry		X		
<i>Gelsemium sempervirens</i>	Florida Jasmine			X	
<i>Haplopappus divaricatus</i>	Scratch daisy				X
<i>Hypericum crux-andreae</i>	St. Peter's wort				X
<i>Hypericum reductum</i>	Atlantic St. John's wort				X
<i>Hypericum gentinoides</i>	Orangeweed				X
<i>Hypericum lissophloeus</i>	Smooth Bark St. John's wort		X		

<i>Hypericum spp.</i>	St. John's wort		X		
<i>Ilex glabra</i>	Gall berry		X		
<i>Ilex myrtifolia</i>	Myrtle leaf holly		X		
<i>Ilex vomitoria</i>	Yaupon		X		
<i>Lachnocaulon anceps</i>	White topped bog buttons				X
					Page 2 of 2
Scientific Name	Common Name	Tree	Shrub	Vine	Herbaceous
<i>Licania michauxii</i>	Gopher apple				X
<i>Lupinus diffusus</i>	Sky-blue lupine				X
<i>Lupinus westianus</i>	Gulf Coast Lupine				X
<i>Magnolia virginiana</i>	Silver bay	X			
<i>Myrica cerifera</i>	Wax myrtle		X		
<i>Opuntia humifusa</i>	Pricklypear cactus				X
<i>Panicum dichotomiflorum</i>	Fall panic grass				X
<i>Panicum hemitomom</i>	Maidencane				X
<i>Paspalum notatum</i>	Bahia grass				X
<i>Persea borbonia</i>	Red Bay	X			
<i>Penstemon multiflorus</i>	Many flowered beardtongue				X
<i>Polygonella gracillis</i>	Wire weed				X
<i>Pinus clausa</i>	Sand Pine	X			
<i>Pinus elliotii</i>	Slash pine	X			
<i>Pinus palustris</i>	Longleaf pine	X			
<i>Pityopsis graminifolia</i>	Golden Aster				X
<i>Polygonella gracilis</i>	Wireweed				X
<i>Quercus geminata</i>	Sand Live Oak	X			
<i>Quercus hemisphaerica</i>	Diamond oak	X			
<i>Quercus incana</i>	Blue jack oak	X			
<i>Quercus laevis</i>	Turkey oak	X			
<i>Quercus virginiana</i>	Live Oak	X			
<i>Rhexia mariana</i>	Pale meadow beauty				X
<i>Rhus copallinum</i>	Sumac		X		
<i>Rubus cuneifolius</i>	Sand blackberry		X		
<i>Schrankia microphylla</i>	Sensitive briar				X
<i>Scoparia dulcis</i>	Sweet Broom				X
<i>Serenoa repens</i>	Saw Palmetto		X		
<i>Smilax sp.</i>	Catbriar			X	
<i>Stylisma sylvatica</i>	Queen's delight				X
<i>Stylisma patens</i>	Coastal plain dawnflower				X
<i>Trichostema dichotomum</i>	Forked blue curls				X
<i>Utricularia floridana</i>	Bladderwort				X
<i>Vaccinium corymbosum</i>	High bush blueberry		X		
<i>Vaccinium myrsinites</i>	Shiny blue berry		X		
<i>Viburnum obovatum</i>	Walter's viburnum		X		
<i>Vitis rotundifolia</i>	Muscadine			X	
<i>Xyris sp.</i>	Yellow-eyed grass				X
<i>Yucca filamentosa</i>	Adam's needle				X

*** Present in previous survey but not observed, ** New observation, * Nuisance Exotic Species

Transect 1

Date 11/4/09

Collector: David Clayton

Wildlife observed: Red wing black bird, titmouse, chipping sparrow
titmouse, chipping sparrow

Community description: Former Sand Pine Plantation


Replanted with LLP

Time: 11:00 Am

Condition, Fair and cool

Scientific Name	Species	Percent Cover	# speices
Aster pilosus	Frost aster	0.5	1
Artemisia campestris	Wormwood	11.6	2
Axonopus furcatus	Big carpet grass	0.6	3
Chrysosoma pauciflosculosa	Woody goldenrod	0.5	4
Cyperus sp.	Sedge	7.6	5
Dichanthelium aciculare	Needle leaf witch grass	0.66	6
Diospyros ebenum	Persimmon	0.16	7
Eupatorium capillifolium	Dog fennel	2	8
Eupatorium mohrii	Mohr's thorough wort	0.5	9
Mollugo verticillata	Indian chickweed	1	10
Opuntia humifusa	Prickly pair cactus	0.167	11
Paspalum notatum	Bahia grass	3.1	12
Pinus clausa	Sand pine	2.3	13
Pinus palustris	Long leaf pine	0.83	14
Quercus hemisphaerica	Diamond oak	0.167	15
Rubus cuneifolius	Sand black berry	1.7	16
Sida rhombifolia	Indian hemp	0.167	17
Tradescantia hirsutiflora	Hairy spiderwort	0.33	18
Yucca filamentosa	Adam's needle	1.6	19
Bare ground	Bare ground	64.1	

Percent Cover



- Bare ground
- Wormwood
- Sedge
- Misc (11 species, .
- Bahia grass
- Sand pine
- Dog fennel
- Sand black berry
- Adam's needle

Transect 1

Date 11/4/09

Collector: David Clayton

Wildlife observed: Red wing black bird, titmouse, chipping sparrow

titmouse, chipping sparrow

Community description: Former Sand Pine Plantation

Time: 11:00 Am

Condition, Fair and cool

Number

LLP

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29

Transect 2

Date 11/4/09

Time: 11:30 Am

Collector: David Clayton

Condition, Fair and cool

Wildlife observed: Titmouse, chipping sparrow

Fuel load: Low

titmouse, chipping sparrow


State Threatened species

Community description: Former Sand Pine Plantation

Wire grass and sandhill species re

Scientific Name	Species	Percent Cover	# speices
Andropogon arctatus	Pinewoods bluestem	1	1
Andropogon glomeratus	Bushy broom sedge	0.17	2
Andropogon virginicus	Broom sedge	0.33	3
Aristida stricta	Wire grass	25.7	4
Axonopus furcatus	Big carpet grass	0.16	5
Cyperus sp.	Sedge	0.16	6
Dichanthelium aciculare	Needle leaf witch grass	0.33	7
Dichanthelium sp.	Witch grass	0.33	8
Diodia teres	Poor Joe	0.33	9
Eragrostis spectabilis	Purple lovegrass	0.167	10
Eugatorium compositifolium	Yankee weed	0.67	11
Gaylussacia dumosa	Dwarf huckleberry	0.33	12
Haplloppus divericatus	Scratch Daisy	0.33	13
Hypericum gentianoides	Orange weed	0.33	14
Liatris tenuifolia	Shortleaf gayfeather	0.67	15
Licania michauxii	Gopher apple	0.33	16
Pinus clausa	Sand pine	0.16	17
Pteridium aquilinum	Bracken	0.15	18
Quercus laevis	Turkey oak	2.2	19
Schizachyrium sp	Little blue stem	1.5	20
Solidago tortifolia	Twisted leaf goldenrod	0.5	21
Stylisma patens	Coastal plain dawkflower	0.16	22
Vaccinium corymbosum	High bush blueberry	1	23
	Bare ground	66	

Percent Cover



- Bare ground
- Wire grass
- Misc (17 species, <: cover)
- Turkey oak
- Little blue stem

Long leaf pine planted

Ht	Condition
1	8
2	8
3	10
4	10
5	12
6	8
7	10
8	10
9	8
10	8
11	8
12	8
13	8
14	8
15	8
16	10

17	8	2
18	10	1
19	6	3
20	6	2
21	6	2
22	6	2
23	8	1
24	6	3
25	10	2
26	10	2
27	8	2
28	10	2
29	8	2
30	8	1
31	8	1
32	8	3
33	6	2
34	8	1
35	8	1
36	6	1
37	6	1
38	6	3
39	6	1
40	5	2
41	10	2
42	8	1
43	10	1
44	10	1
45	6	2
46	8	1
47	8	1
48	6	1
49	6	2
50	8	3


399 87
7.98 1.74

Transect 3
Date 11/4/09
Collector: David Clayton
Wildlife observed: Crow
Community description: Sandhill with oaks thinned
Time: 1:00 Pm
Condition, Fair and cool
Fuel load: Moderate
Greenhead branch

Scientific Name	Species	Percent Cover	# speices
Andropogon arctatus	Pinewoods bluestem	3.3	1
Andropogon virginicus	Broom sedge	0.167	2
Aristida stricta	Wire grass	39	3
Chrysoma pauciflosculosa	Woody goldenrod	1.7	4
Diospyros virginiana	Persimon	0.167	5
Gaylussacia dumosa	Dwarf huckleberry	1	6
Persea borbonia	Red bay	1	7
Pinus palustris	Long leaf pine	0.33	8
Polygonella gracilis	Wire weed	0.167	9
Opuntia humifusa	Prickly pair cactus	0.167	10
Quercus incana	Blue jack oak	0.167	11
Quercus laevis	Turkey oak	3.5	12
Serenoa repens	Saw palmetto	5.3	13
Vaccinium darrowii	Darrow's blueberry	0.167	14
Vaccinium arboreum	Sparkle berry	0.167	15
	Bare ground	44	



■ Bare ground



Ht	Condition
1	12
2	10
3	10
4	6
5	24
6	10
7	24
8	20
9	6
10	20
11	8
12	12
13	10
14	20
15	20
16	6
17	20
18	20
19	10
20	12

21	10	1
22	24	1
23	24	1
24	8	1
25	24	1
26	20	1
27	36	3
28	36	4
29	24	2
30	24	1
31	12	1
32	8	3
33	6	2
34	24	1
35	36	1
36	36	1
37	12	1
38	12	1

656	1.342105263
17.26315789	51

Transect 4

Date 11/5/09

Collector: David Clayton

Wildlife observed: None

Community description: Sand pine plantation removed

Time: 10:30 Pm

Condition, Fair and cool


Fuel load: low

East side Dykes Mill Pond and Gre

Scientific Name	Species	Percent Cover	# species
Aeschynomene americana	Aeschynomene	0.167	1
Andropogon virginicus	Broom sedge	0.33	2
Aristida stricta	Wire grass	12.3	3
Bulbostylis ciliatifolia	Capillary hair sedge	0.167	4
Chrysoma pauciflosculosa	Woody goldenrod	2.3	5
Chrysopsis lanuginosa	Lynn Haven goldenaster	1.5	6
Conyza canadensis	Canadian horseweed	1.6	7
Cyperus sp.	Sedge	0.83	8
Dichaenanthelium sp.	Witch grass	1	9
Diodia teres	Poor Joe	2	10
Diospyros virginiana	Persimon	0.5	11
Eremochloa ophiuroides	Centpede grass	4	12
Eupatorium capillifolium	Dog fennel	13	13
Eupatorium mohrli	Mohr's thorough wort	1.8	14
Galactia sp.	Milk pea	0.167	15
Hypericum gentianoides	Orange weed	1.6	16
Ilex vomitoria	Yaupon	0.5	17
Pinus palustris	Long leaf pine	0.83	18

Opuntia humifusa	Prickly pair cactus	1.67	19
Quercus hemaespherica	Diamond Oak	0.33	20
Rubus cuneifolius	Sand black berry	4	21
Schizachyrium sp	Little blue stem	1.3	22
Vaccinium arboreum	Sparkle berry	0.16	23
	Bare ground	48	24
			25
			26

Percent Cover




- Bare ground
- Dog fennel
- Wire grass
- Centipede grass
- Sand black berry
- Misc (10 species, 1% co)
- Woody goldenrod
- Poor Joe
- Mohr's thorough wort
- Prickly pair cactus
- Canadian horseweed
- Orange weed
- Lynn Haven goldenastei
- Little blue stem
- Witch grass

LLP

Height

1	8	1
2	8	1
3	8	1
4	10	1
5	8	1
6	8	1
7	8	1
8	12	1
9	12	1
10	8	1
11	10	1
12	8	1
13	8	1
14	6	1
15	6	2
16	8	2
17	8	2
18	8	1
19	8	1
20	8	3
21	8	2
22	8	1
23	8	3
24	8	1

8.3333333333 1.3333333333



Transect 5

Date 11/5/09

Collector: David Clayton

Wildlife observed: Wren, chipping sparrow

Community description: Sandhill

Time: 8:50 am


Condition, Fair and cool


Fuel load: moderate due to herbic

Scientific Name	Species	Percent Cover	# speices
<i>Andropogon arctatus</i>	Pinewoods bluestem	1.3	1
<i>Andropogon virginicus</i>	Broom sedge	0.33	2
<i>Aristida stricta</i>	Wire grass	42.8	3
<i>Asimina angustifolia</i>	Slimleaf paw paw	0.167	4
<i>Aster pilosus</i>	Frost aster	0.167	5
<i>Balduna angustifolia</i>	Coastalplain honeycomb head	1.5	6
<i>Baptisia lanceolata</i>	Gopherweed	0.167	7
<i>Bulbostylis ciliatfolia</i>	Capillary hair sedge	0.167	8
<i>Chrysoma pauciflosculosa</i>	Woody goldenrod	2.67	9
<i>Crataegus michauxii</i>	Michaux's hawthorn	0.33	10
<i>Dichanthelium sp.</i>	Witch grass	0.167	11
<i>Diospyros virginiana</i>	Persimon	2	12
<i>Eriogonum tomentosum</i>	buckwheat	0.167	13
<i>Gelsemium sempervirens</i>	Yellow jessamine	0.167	14
<i>Hypericum gentianoides</i>	Orange weed	0.5	15
<i>Ilex vomitoria</i>	Yaupon	0.167	16
<i>Licania michauxii</i>	Gopher apple	1.167	17
<i>Penstemon multiflorus</i>	Manyflowered beardtongue	0.167	18
<i>Pityopsis graminifolia</i>	Narrow leaf silkgrass	0.167	19
<i>Polygonella gracilis</i>	Wire weed	5.8	20
<i>Pteridium aquilinum</i>	Bracken	1.5	21
<i>Opuntia humifusa</i>	Prickly pair cactus	0.66	22
<i>Quercus incana</i>	Blue jack oak	0.167	23

Scleria sp.	Nutrush	0.167	24
Smilax sp.	Smilax	0.83	25
Vaccinium corymbosum	High bush blueberry	0.83	26
Vaccinium myrsinites	Shiny blueberry	1.3	27
Yucca filimentosa	Adam's needle	0.33	28
	Bare ground	33.5	

Percent Cover





Transect 6

Date 11/3/09

Collector: David Clayton

Wildlife observed: titmouse

Community description: Wet flatwoods

Time: 10:00 am


Condition, Fair and cool

Fuel load: low


Scientific Name	Species	Percent Cover	# speices
Aristida stricta	Wire grass	3.8	1
Citronia monophylla	Black titi	4.6	2
Cyrtilla racemiflora	Red titi	1	3
Ilex glabra	Gall berry	1.3	4
Ilex myrtifolia	Myrtle-leaved holly	2.6	5
Lachnanthes caroliniana	Red root	0.16	6
Lyonia lucida	Fetter bush	16.1	7
Osmanthus americanus	Wild olive	0.16	8

Persea palustris	Swamp bay	0.66	9
Rhynchospora microcephala	Bunched beaksedge	3.2	10
Vaccinium corymbosum	Highbush blueberry	0.167	11
	Bare ground	66.5	

Percent Cover



- Bare gr
- Fetter l
- Black ti
- Wire gr
- Bunche
- Myrtle-
- Gall bei
- Misc sp
- Red titi



Transect 7

Date 11/2/09

Collector: David Clayton

Wildlife observed:none

Community description: Wet flatwoods

Time: 10:00 am


Condition, Fair and cool

Fuel load:low

Scientific Name	Species	Percent Cover	# speices
<i>Andropogon glomeratus</i>	Bushy bluestem	0.67	1
<i>Aristida stricta</i>	Wire grass	2.8	2
<i>Cliftonia monophylla</i>	Black titi	18.3	3
<i>Cyrilla racemiflora</i>	Red titi	5.1	4
<i>Eupatorium capillifolium</i>	Dog fennel	0.5	5
<i>Gaylussacia dumosa</i>	Dwarf blueberry	0.8	6
<i>Ilex glabra</i>	Gall berry	0.5	7
<i>Ilex myrtifolia</i>	Myrtle-leaved holly	0.16	8

<i>Lachnanthes caroliniana</i>	Red root	5.5	9
<i>Leucothoe racemosa</i>	Swamp dog hobble	0.167	10
<i>Lyonia lucida</i>	Fetter bush	1.7	11
<i>Persea palustris</i>	Swamp bay	0.167	12
<i>Rhynchospora microcephala</i>	Bunched beaksedge	0.167	13
<i>Vaccinium corymbosum</i>	Highbush blueberry	1.5	14
	Bare ground	61.1	

Percent Cover





Transect 8

Date 11/2/09

Time: 11:30 am

Collector: David Clayton

Condition, Fair and cool

Wildlife observed: Red shouldered hawk, titmouse, red bellied woodpecker

Community description: Wet flatwoods, slash pine thinned

Fuel load: low

Scientific Name	Species	Percent Cover	# speices
<i>Andropogon glomeratus</i>	Bushy bluestem	0.167	1
<i>Carex verrucosa</i>	Warty Sedge	0.5	2
<i>Centella asiatica</i>	Centella	0.8	3
<i>Chamaecrista nictitans</i>	Partridge berry	1.1	4
<i>Cliftonia monophylla</i>	Black titi	5.1	5
<i>Cyrtia racemiflora</i>	Red titi	0.167	6
<i>Dichantheilium sp.</i>	Witch grass	3.16	7
<i>Diodia teres</i>	Poor joe	2.3	8
<i>Eupatorium capillifolium</i>	Dog fennel	1.5	9
<i>Gelsemium sempervirens</i>	Jessamine	0.33	10
<i>Hyper sp.</i>	St. John's wort	1	11
<i>Ilex glabra</i>	Gall berry	0.5	12
<i>Ilex vomitoria</i>	Yaupon	1	13
<i>Lochneranthes caroliniana</i>	Red root	6.5	14
<i>Leucothoe racemosa</i>	Swamp dog hobble	0.167	15
<i>Ludwigia decurrens</i>	Seedbox	0.5	16
<i>Lycopus virginicus</i>	Water horehound	0.3	17
<i>Myrica cerifera</i>	Wax myrtle	1.8	18
<i>Panicum sp.</i>	Panic grass	8	19
<i>Persea palustris</i>	Swamp bay	0.8	20
<i>Pinus palustris</i>	Longleaf pine	0.1	21
<i>Polygala lutea</i>	Candy weed	0.16	22
<i>Rhexia alifanous</i>	Savannah meadow beauty	0.06	23
<i>Rhynchospora microcephala</i>	Bunched beaksedge	0.167	24
<i>Rubus cuneifolius</i>	Sand black berry	1.3	25
<i>Smilax laurifolia</i>	Cat briar	0.167	26
<i>Vaccinium corymbosum</i>	Highbush blueberry	0.3	27

Vitis rotundifolia

Muscadine grape


Bare ground

9.5

52

28

Species







Transect 9

Date 11/6/09

Collector: David Clayton

Wildlife observed: Great white egret, little blue heron, alligator

Community description: Marsh


Time: 9:30 am

Condition: Fair and cool

Fuel load: N/A

Scientific Name	Common Name	Percent Cove	# species
Drosera intermedia	Water sundew (Fl threate	0.16	1
Eleocharis cellulosa	Eleocharis	3.3	2
Lachnanthes caroliniana	Red root	0.5	3
Nuphar advena subsp. Orbicul	Spatterdock	4.7	4
Nymphaea odorata	Fragrant water lily	35.5	5
Rhynchospora inundata	Horned beaksedge	6.8	6
Utricularia floridana	Florida bladderwort	1.2	7
Websteria confervoides	Websteria	7.5	8
Xyris sp	Yellow-eyed grass	2.3	9
	Open water	38	10

Percent Cover







Quadrat Transect 1

Species	Cover	Species	Cover	Species	Percent Cover
1 Art Cam	40 Art Cam	40	Bare ground	64.1	
Cyp sp	10 Art Cam	80	Wormwood	11.6	
Bare ground	50 Art Cam	75	Sedge	7.6	
2 Art Cam	80 Art Cam	30	Misc (11 species, < 1% cover)	4.6	
Pin paul	10 Art Cam	10	Bahia grass	3.1	
Bare ground	10 Art Cam	20	Sand pine	2.3	
3 Art Cam	75 Art Cam	45	Dog fennel	2	
Bare ground	25 Art Cam	5	Sand black berry	1.7	
4 Bare ground	100 Art Cam	45	Adam's needle	1.6	
5 Moll vert	30 Ast pil	15	Indian chickweed	1	
Bare ground	70 Axon furc	20	Long leaf pine	0.83	
6 Bare ground	100 Bare ground	50	Needle leaf witch grass	0.66	
7 Bare ground	100 Bare ground	10	Big carpet grass	0.6	
8 Art Cam	30 Bare ground	25	Frost aster	0.5	
Bare ground	70 Bare ground	100	Woody goldenrod	0.5	
9 Eup cap	30 Bare ground	70	Mohr's thorough wort	0.5	
Trad hir	5 Bare ground	100	Hairy spiderwort	0.33	
Bare ground	65 Bare ground	100	Prickly pair cactus	0.167	
10 Dichan acuc	15 Bare ground	70	Diamond oak	0.167	
Cyp sp	50 Bare ground	65	Indian hemp	0.167	
Bare ground	35 Bare ground	35	Persimon	0.16	
11 Art Cam	10 Bare ground	65			
n paul	5 Bare ground	30			
ib cun	5 Bare ground	40			
yp sp	10 Bare ground	45			
ad hir	5 Bare ground	15			
are ground	65 Bare ground	30			
ib cun	15 Bare ground	40			
ip cap	20 Bare ground	45			
yp sp	15 Bare ground	55			
t Cam	20 Bare ground	85			
are ground	30 Bare ground	100			

< 1% cover)

14	Art Cam	50	Bare ground	90		
	Rub cun	10	Bare ground	85		
	Bare ground	40	Bare ground	100		
	Art pil	15	Bare ground	90		
	Chry pau	15	Bare ground	95		
	Eup moh	15	Bare ground	55		
	Dios eben	5	Bare ground	95		
	Cyp sp	5	Bare ground	50		
15	Bare ground	45	Bare ground	90	64.16667	64.16667
	pas pal	80	Chry pau	15		0.5
	Rub cun	5	Cyp sp	10		
	Bare ground	15	Cyp sp	50		
16	Pin clau	30	Cyp sp	10		
	Cyp sp	40	Cyp sp	15		
	Bare ground	30	Cyp sp	10		
17	Pin clau	30	Cyp sp	5		
	Cyp sp	10	Cyp sp	40		
	Axon furc	20	Cyp sp	10		
	Bare ground	40	Cyp sp	35		
18	Art Cam	45	Cyp sp	10		
	Eup cap	10	Cyp sp	5		
	Bare ground	45	Cyp sp	5		
19	Cyp sp	35	Cyp sp	20		
	Art Cam	5	Cyp sp	5	7.666667	
	Rub cun	5	Dichan acuc	15		
	Bare ground	55	Dichan acuc	5	0.666667	
20	sid rhom	5	Dios eben	5	0.166667	
	Cyp sp	10	Eup cap	30		
	Bare ground	85	Eup cap	20		
21	Bare ground	100	Eup cap	10		2
22	Opun hum	5	Eup moh	15		0.5
	Cyp sp	5	Moll vert	30		1
	Bare ground	90	Opun hum	5	0.166667	
23	Rub cun	15	pas pal	80		

Bare ground	85 pas pal	15	3.166667
24 Bare ground	100 Pin clau	30	
25 Pin clau	10 Pin clau	30	
Bare ground	90 Pin clau	10	2.333333
26 Cyp sp	5 Pin paul	10	
Bare ground	95 Pin paul	5	
27 Art Cam	45 Pin paul	10	0.833333
Bare ground	55 Quer hem	5),.166667
28 Rub cun	5 Rub cun	5	
Bare ground	95 Rub cun	15	
29 Pin paul	10 Rub cun	5	
pas pal	15 Rub cun	5	
Quer hem	5 Rub cun	15	1.666667
Cyp sp	20 Rub cun	5	
Bare ground	50 sid rhom	5	0.1667
30 Cyp sp	5 Trad hir	5	
Dichan acuc	5 Trad hir	5	0.333333
Bare ground	90 Yuc fill	50	1.666667

3000 99.7667

Size
inches

8	1
8	1
6	2
8	1
8	1
6	2
6	2
6	2

10	1
10	1
6	2
8	2
12	1
10	1
8	2
6	2
10	2
8	2
8	2
8	1
10	1
8	2
6	3
8	2
8	2
6	3
10	2
12	1
10	1
238	48
8.2069	1.655172414

generating

Quadrat **Transect 2**
Species **Cover** **Species** **Cover**

1	ari str	90	And arct	35			
	Bare ground	10	And arct	30	1		
2	Gay dum	5	And glom	5	0.166667	Brachen	
	And arct	35	And glom	5		Big carpet grass	
	ari str	15	And vir	10	0.333333	Sedge	
	Bare ground	45	And vir	5		Sand pine	
3	Styl pat	5	ari str	90		Coastal plain dafflower	
	ari str	20	ari str	15		Purple lovegrass	
	Bare ground	75	ari str	20		Bushy broom sedge	
4	Dich aci	5	ari str	20		Broom sedge	
	Bare ground	95	ari str	70		Needle leaf witch grass	
5	Bare ground	100	ari str	60		Witch grass	
6	Hap div	10	ari str	25		Poor Joe	
	Sol tor	15	ari str	5		Dwarf huckleberry	
	pin cla	5	ari str	50		Scratch Daisy	
	And vir	10	ari str	80		Orange weed	
	Eup comp	20	ari str	5		Gopher apple	
	Cyperus sp	5	ari str	5		Twisted leaf goldenrod	
	Axo fur	5	ari str	35		Yankee weed	
	Bare ground	30	ari str	70		Speices	
7	ari str	20	ari str	50		Bare ground	Percent Cover
	Bare ground	80	ari str	30		Wire grass	66
8	Quer lae	5	ari str	25		Misc (17 species, <1% cover)	25.7
	ari str	70	ari str	20		Turkey oak	2.2
	Bare ground	25	ari str	30		Little blue stem	1.5
9	ari str	60	ari str	30	770	Pinewoods bluestem	1
	Bare ground	40	ari str	35	25.666667	High bush blueberry	1
10	ari str	25	Axo fur	5	0.166667		
	Bare ground	75	Bare ground	10			
11	Pter aqu	5	Bare ground	45			
	Dic sp	10	Bare ground	75			
	ari str	5	Bare ground	95			
	Quer lae	10	Bare ground	100			
	Bare ground	70	Bare ground	30			

1%

12 ari str	50 Bare ground	80	
Bare ground	50 Bare ground	25	
13 Bare ground	100 Bare ground	40	
14 ari str	80 Bare ground	75	
Bare ground	20 Bare ground	70	
15 sch sp	15 Bare ground	50	
ari str	5 Bare ground	100	
Gay dum	5 Bare ground	20	
Bare ground	75 Bare ground	75	
16 Lic mich	5 Bare ground	65	
And arct	30 Bare ground	80	
Bare ground	65 Bare ground	65	
17 And glom	5 Bare ground	30	
Quer lae	5 Bare ground	20	
ari str	5 Bare ground	70	
Lic mich	5 Bare ground	70	
Bare ground	80 Bare ground	40	
18 ari str	35 Bare ground	85	
Bare ground	65 Bare ground	90	
19 ari str	70 Bare ground	65	
Bare ground	30 Bare ground	65	
20 Vac cor	30 Bare ground	65	
ari str	50 Bare ground	60	62
Bare ground	20 Bare ground	100	1860
21 ari str	30 Cyperus sp	5	0.166667
Bare ground	70 Dic sp	10	0.333333
22 Era spec	5 Dich aci	5	
ari str	25 Dich aci	5	0.33
Bare ground	70 Diodia teres	10	0.33
23 Quer lae	40 Era spec	5	0.166667
ari str	20 Eup comp	20	0.666667
Bare ground	40 Gay dum	5	
24 Quer lae	5 Gay dum	5	0.33
And glom	5 Hap div	10	0.33

Dich aci	5 Hyp gen	5	
Bare ground	85 Hyp gen	5	0.33
25 And vir	5 Lia ten	5	0.16
Lia ten	5 Lic mich	5	
Bare ground	90 Lic mich	5	0.33
26 ari str	30 pin cla	5	0.16
Hyp gen	5 Pter aqu	5	0.15
Bare ground	65 Quer lae	5	
27 ari str	30 Quer lae	10	
Hyp gen	5 Quer lae	5	
Bare ground	65 Quer lae	40	2.166667
28 ari str	35 Quer lae	5	65
Bare ground	65 sch sp	15	
29 sch sp	30 sch sp	30	1.5
Diodia teres	10 Sol tor	15	0.5
Bare ground	60 Styl pat	5	0.16
30 Bare ground	100 Vac cor	30	1

3000

Quadrat Transect 3

Species	Cover	Species	Cover	Species	Cover
1 Ari str	70	Species	Cover	Species	cover
Pin paul	10	And arc	5	Blue jack oak	
bar grnd	20	And arc	5	Broom sedge	
2 Quer lae	25	And arc	10	Darrow's blueberry	
Ari str	20	And arc	5	Persimon	
bar grnd	55	And arc	55	Prickly pair cactus	
3 And vir	5	And arc	20	Sparkle berry	
And arc	5	And vir	5	Wire weed	
Chry pau	5	Ari str	70	Long leaf pine	
gay dumosa	5	Ari str	20		
bar grnd	80	Ari str	10		
4 And arc	5	Ari str	5	Species	Cover
Ari str	10	Ari str	20	Bare ground	44
Quer lae	10	Ari str	30	Wire grass	39
gay dumosa	5	Ari str	30	Saw palmetto	5.3
bar grnd	70	Ari str	5	Turkey oak	3.5
5 Ari str	5	Ari str	25	Pinewoods bluestem	3.3
bar grnd	95	Ari str	20	Woody goldenrod	1.7
6 Ari str	20	Ari str	30	Misc (8 species, < 1% cover)	1.5
bar grnd	80	Ari str	30	Dwarf huckleberry	1
7 Ari str	30	Ari str	55	Red bay	1
pol gra	5	Ari str	95		
And arc	10	Ari str	80		
Chry pau	10	Ari str	30		
bar grnd	45	Ari str	40		
8 Ser rep	60	Ari str	55		
Ari str	30	Ari str	30		
bar grnd	10	Ari str	95		
9 Ser rep	95	Ari str	50		
bar grnd	5	Ari str	70		

10	Quer lae	20	Ari str	35	
	Chry pau	10	Ari str	35	
	Ari str	5	Ari str	40	38.66667
	bar grnd	65	Ari str	65	1160
11	Ari str	25	Ari str	90	#NAME?
	bar grnd	75	bar grnd	20	
12	Quer lae	10	bar grnd	55	
	Opu hum	5	bar grnd	80	
	gay dumosa	10	bar grnd	70	
	Ari str	20	bar grnd	95	
	bar grnd	55	bar grnd	80	
13	Ari str	30	bar grnd	45	
	And arc	5	bar grnd	10	
	Chry pau	5	bar grnd	5	
	gay dumosa	5	bar grnd	65	
	bar grnd	55	bar grnd	75	
14	Ari str	30	bar grnd	55	
	Chry pau	5	bar grnd	55	
	gay dumosa	5	bar grnd	60	
	bar grnd	60	bar grnd	40	
15	Ari str	55	bar grnd	5	
	Quer lae	5	bar grnd	20	
	bar grnd	40	bar grnd	65	
16	Ari str	95	bar grnd	25	
	bar grnd	5	bar grnd	35	
17	Ari str	80	bar grnd	65	
	bar grnd	20	bar grnd	45	
18	Quer incan	5	bar grnd	5	
	Ari str	30	bar grnd	35	
	bar grnd	65	bar grnd	10	
19	Per bor	30	bar grnd	55	
	Ari str	40	bar grnd	65	
	Dios vir	5	bar grnd	60	44
	bar grnd	25	bar grnd	15	1320

20 Ari str	55 bar grnd	5	1320
Chry pau	5 Chry pau	5	
vac arbo	5 Chry pau	10	
bar grnd	35 Chry pau	10	
21 Ari str	30 Chry pau	5	
Chry pau	5 Chry pau	5	
bar grnd	65 Chry pau	5	
22 And arc	55 Chry pau	5	50
Quer lae	5 Chry pau	5	1.666667
bar grnd	45 Dios vir	5	0.167
23 Ari str	95 gay dumosa	5	
bar grnd	5 gay dumosa	5	
24 Ari str	50 gay dumosa	10	
Chry pau	5 gay dumosa	5	
Ser rep	5 gay dumosa	5	1
Quer lae	5 Opu hum	5	0.167
bar grnd	35 Per bor	30	1
25 Ari str	70 Pin paul	10	0.333333
Quer lae	15 pol gra	5	0.167
bar grnd	10 Quer incan	5	0.167
26 Ari str	35 Quer lae	25	
Quer lae	10 Quer lae	10	
bar grnd	55 Quer lae	20	
27 Ari str	35 Quer lae	10	
bar grnd	65 Quer lae	5	
28 Ari str	40 Quer lae	5	
bar grnd	60 Quer lae	5	3.5
29 Ari str	65 Quer lae	15	
And arc	20 Quer lae	10	105
bar grnd	15 Ser rep	60	
30 vac dar	5 Ser rep	95	5.333333
Ari str	90 Ser rep	5	160
bar grnd	5 vac arbo	5	0.166667
	vac dar	5	0.1667

3000

en Head

Quadrat Transect 4

1 Species	Cover	Species	Cover		Species
Ari str	40	Aesh amer	5	0.167	
Eup mohr	20	And vir	10	0.333333	
Bare grnd	40	Ari str	40		
2 Ari str	40	Ari str	40		
Bare grnd	60	Ari str	5		
3 pin pal	5	Ari str	20		
Eup cap	20	Ari str	20		
Sch sp	10	Ari str	30		
Bare grnd	65	Ari str	10		
4 Chry lan	15	Ari str	15		
Dios vir	5	Ari str	10		
Dich sp	5	Ari str	15		
Ari str	5	Ari str	40		
Bare grnd	70	Ari str	15		
5 Eup cap	50	Ari str	15		
Conz can	20	Ari str	5		
Bare grnd	30	Ari str	10		
6 Ari str	20	Ari str	20		
Eup cap	40	Ari str	25		
Chry lan	15	Ari str	25	370	Aristida stricta
Bare grnd	25	Ari str	10	12.333333	
7 Ari str	20	Bare grnd	40		
Eup cap	40	Bare grnd	60		
Chry lan	15	Bare grnd	65		
Bare grnd	25	Bare grnd	70		

Species

- Bare ground
- Dog fennel
- Wire grass
- Centipede grass
- Sand black berry
- Misc (10 species, 1% cover)
- Woody goldenrod

8 Ari str	30 Bare grnd	30		Poor Joe
Eup mohr	10 Bare grnd	25		Mohr's thorough wort
Bare grnd	60 Bare grnd	25		Prickly pair cactus
9 Pin pal	5 Bare grnd	60		Canadian horseweed
Rub arg	60 Bare grnd	30		Orange weed
Cyp sp	5 Bare grnd	40		Lynn Haven goldenaster
Bare grnd	30 Bare grnd	65		Little blue stem
10 Sch sp	20 Bare grnd	70		Witch grass
Pin pal	10 Bare grnd	20		
Conz can	10 Bare grnd	15		
Hyp gen	10 Bare grnd	70		
Rub arg	10 Bare grnd	70		
Bare grnd	40 Bare grnd	10		
11 Galac sp	5 Bare grnd	75		
Rub arg	20 Bare grnd	55		
Ari str	10 Bare grnd	70		
Bare grnd	65 Bare grnd	85		
12 Dios vir	10 Bare grnd	100		Aeschynomene americana
Ari str	15 Bare grnd	30		Aeschynomene
Bare grnd	5 Bare grnd	35		Broom sedge
Cyp sp	70 Bare grnd	20		Capillary hair sedge
Bare grnd	25 Bare grnd	40		Sedge
13 Opun hum	50 Bare grnd	30		Persimon
Dio ter	5 Bare grnd	40		Milk pea
Ten	20 Bare grnd	25	1440	Yaupon
Bare grnd	20 Bare grnd	70	48	Long leaf pine
14 Ere oph	40 Chry lan	15		Diamond Oak
Eup cap	15 Chry lan	15		Sparkle berry
Rub arg	10 Chry lan	15	1.5	
Ari str	15 Conz can	20		
Bare grnd	15 Conz can	10		
15 Ere oph	15 Conz can	5	1.166667	
Hyp gen	70 Cry pau	40		
Bare grnd	15 Cry pau	20		
16 Ari str				

Dich sp	10 Cry pau	10	2.333333
Aesh amer	5 Cyp sp	5	
Bare grnd	70 Cyp sp	5	
17 Ari str	40 Cyp sp	5	
Ere oph	10 Cyp sp	5	
Eup cap	30 Cyp sp	5	0.833333
Dich sp	10 Dich sp	5	
Bare grnd	10 Dich sp	10	
18 Cyp sp	5 Dich sp	10	
Ari str	15 Dich sp	5	1
Dich sp	5 Dio ter	50	
Bare grnd	75 dio ter	10	2
19 Opun hum	20 Dios vir	5	
Conz can	5 Dios vir	10	0.5
pin pal	5 Ere oph	20	
Ari str	15 Ere oph	15	
Bare grnd	55 Ere oph	10	
20 Cyp sp	5 Ere oph	15	
Hyp gen	20 Ere oph	40	
Ari str	5 Ere oph	10	
Bare grnd	70 Ere oph	10	4
21 Eup cap	5 Eup cap	20	
Ari str	10 Eup cap	50	
Bare grnd	85 Eup cap	40	
22 Bare grnd	100 Eup cap	40	
23 Cry pau	40 Eup cap	40	
Eup mohr	25 Eup cap	30	
Rub arg	5 Eup cap	5	
Bare grnd	30 Eup cap	40	
24 Ere oph	15 Eup cap	30	
Cry pau	20 Eup cap	15	
Ari str	20 Eup cap	40	13
dio ter	10 Eup cap	40	390
Bare grnd	35 Eup mohr	20	

Transect 5					
Quadrat	Species	Cover	Species	Cover	Percent Cover
1	Vac myr	15	And arc	5	42.8
	Ari str	25	And arc	5	33.5
	Bar grnd	60	And arc	10	5.8
2	Ari str	25	And arc	5	5.8
	Bar grnd	75	And arc	5	2.67
3	Ari str	45	And arc	10	2
	Bar grnd	55	And vir	10	1.5
	pol gra	5	Ari str	0.333333	1.5
	bauld ang	10	Ari str	25	1.5
	bulb cil	5	Ari str	45	1.3
	Ari str	20	Ari str	20	1.3
	Bar grnd	60	Ari str	35	1.167
5	Vac cor	25	Ari str	60	0.83
	Ari str	35	Ari str	40	0.83
	Dich sp	5	Ari str	30	0.66
	pol gra	5	Ari str	40	0.5
	Bar grnd	30	Ari str	45	0.33
6	Vac myr	5	Ari str	50	0.33
	bauld ang	20	Ari str	50	0.33
	And arc	5	Ari str	60	0.167
	Ari str	60	Ari str	20	0.167
	Bar grnd	5	Ari str	45	0.167
7	pter aqu	10	Ari str	45	0.167
	Ari str	40	Ari str	20	0.167
	Smi sp.	5	Ari str	75	0.167
	Bar grnd	40	Ari str	25	0.167
	Crat mich	5	Ari str	25	0.167
8	Ari str	30	Ari str	60	0.167
	pol gra	10	Ari str	70	0.167
	lic mich	5	Ari str	55	0.167
	Ilex vom	5	Ari str	20	5.814

dios vir	10 Ari str	50	
pter aqu	5 Ari str	80	
Smi sp.	5 Ari str	30	
Bar grnd	30 Ari str	50	1285
9 Yucca filimentosa	10 Ari str	90	42.833333
dios vir	5 Asim ang	5	0.166667
pol gra	5 Ast pol	5	0.1667
bauld ang	5 Bapt lan	5	0.1667
	nd	60	
	nd	75	
	nd	55	
	nd	60	
	nd	30	
	nd	5	
	nd	40	
	nd	30	
	nd	30	
	nd	35	
	nd	15	
	nd	25	
	nd	25	
	nd	60	
	nd	10	
	nd	20	
	nd	5	
	nd	5	
	nd	40	
	nd	65	
	nd	50	
	nd	15	
	nd	5	
	nd	45	
	nd	75	
	nd	40	

- Wire grass
- Bare ground
- Wire weed
- Misc (19 species, < 1% cover)
- Woody goldenrod
- Persimon
- Coastalplain honeycomb head
- Brachen
- Pinewoods bluestem
- Shiny blueberry
- Gopher apple

15 Ari str	45 Bare grnd	5	
Chry pau	20 Bare grnd	60	
pol gra	10 Bare grnd	15	1005
And arc	5 bauld ang	5	33.5
dios vir	5 bauld ang	10	
smil sp	5 bauld ang	20	
Bare grnd	10 bauld ang	5	
16 Opun hum	20 bauld ang	5	1.5
Ari str	45 bulb cil	5	0.1667
Vac myr	10 Chry pau	5	
dios vir	5 Chry pau	20	
Bare grnd	20 Chry pau	55	2.666667
17 Chry pau	55 Crat mich	5	
Ari str	20 Crat mich	5	0.333333
pter aqu	10 Dich sp	5	0.166667
Ast pol	5 dios vir	10	
pol gra	5 dios vir	5	
Bare grnd	5 dios vir	5	
18 Scler	5 dios vir	5	
pol gra	5 dios vir	5	
Ari str	75 dios vir	5	
Bapt lan	5 dios vir	5	
Asim ang	5 dios vir	5	
Bare grnd	5 erio tomen	20	2
19 pter aqu	10 gel sem	5	0.1667
Ari str	25 Hyper gen	5	
quer inc	5 Hyper gen	5	
dios vir	5 Hyper gen	5	0.5
pen mult	5 llex vom	5	0.1667
lic mich	5 lic mich	5	
Vac myr	10 lic mich	5	
Bare grnd	40 lic mich	10	
20 erio tomen	5 lic mich	5	

Ari str	25 lic mich	5	
pol gra	5 lic mich	5	1.166667
Bare grnd	65 Opun hum	20	0.666667
21 And vir	10 pen mult	5	0.1667
pol gra	30 pity gra	5	0.1667
Crat mich	5 pol gra	5	
smil sp	5 pol gra	5	
Bare grnd	50 pol gra	10	
22 Ari str	60 pol gra	5	
pter aqu	10 pol gra	10	
And arc	5 pol gra	20	
pol gra	10 pol gra	15	
Bare grnd	15 pol gra	10	
23 bauld ang	5 pol gra	10	
pol gra	10 pol gra	5	
Hyper gen	5 pol gra	5	
gel sem	5 pol gra	5	
Ari str	70 pol gra	30	
Bare grnd	5 pol gra	10	
24 Ari str	55 pol gra	10	
Bare grnd	45 pol gra	10	175
25 Ari str	20 pol gra	10	5.833333
Hyper gen	5 pter aqu	10	
Bare grnd	75 pter aqu	5	
26 Ari str	50 pter aqu	10	
And arc	10 pter aqu	10	
Bare grnd	40 pter aqu	10	
27 Ari str	80 pter aqu	10	1.5
pol gra	10 quer inc	3	
dios vir	5 quer inc	5	0.266667
Bare grnd	5 Scler	5	0.1667
28 Ari str	30 smi sp	2	
dios vir	5 smi sp	5	
Hyper gen	5 Smi sp.	5	

Bare grnd	60 Smi sp.	5	
29 Ari str	50 smil sp	5	
dios vir	20 smil sp	5	0.833333
pol gra	10 Vac cor	25	0.83
lic mich	5 Vac myr	15	
Bare grnd	15 Vac myr	5	
30 Ari str	90 Vac myr	10	
lic mich	5 Vac myr	10	1.333333
Bare grnd	5 Yucca filimentosa	10	0.333333

3000

Quadrat	Transect 6	Cover	Species	Cover	Species	Percent Cover
1	Lyonia luc	5	Ari str	5	Bare ground	66.5
	Cli mon	5	Ari str	5	Fetter bush	16.1
	Bare grnd	90	Ari str	5	Black titi	4.6
2	Lyonia luc	10	Ari str	5	Wire grass	3.8
	Cli mon	3	Ari str	10	Bunched beaksedge	3.2
	Bare grnd	87	Ari str	10	Myrtle-leaved holly	2.6
3	Lyonia luc	15	Ari str	25	Gall berry	1.3
	Ari str	5	Ari str	5	Misc species (4, < 1% cover)	1.2
	Bare grnd	80	Ari str	10	Red titi	1
4	Lyonia luc	20	Ari str	10	Swamp bay	0.66
	Cli mon	15	Ari str	10	Highbush blueberry	0.167
	Bare grnd	65	Ari str	10	Red root	0.16
5	cyr rac	10	Ari str	5	Wild olive	0.16
	Per pal	20	Bare grnd	90		1.147
		30	Bare grnd	87		
		40	Bare grnd	80		

ound	20 Bare grnd	65	
bush	80 Bare grnd	40	
iti	100 Bare grnd	80	
'ass	5 Bare grnd	100	
nd beaksedge	15 Bare grnd	45	
-leaved holly	35 Bare grnd	35	
rty	45 Bare grnd	70	
pecies (4, < 1% cover)	5 Bare grnd	50	
	20 Bare grnd	60	
	20 Bare grnd	60	
	20 Bare grnd	60	
	20 Bare grnd	50	
	35 Bare grnd	95	
	15 Bare grnd	90	
	15 Bare grnd	60	
	70 Bare grnd	85	
	25 Bare grnd	95	
	20 Bare grnd	85	
	5 Bare grnd	70	
Ari str	50 Bare grnd	65	
Bare grnd	10 Bare grnd	20	
12 Cli mon	25 Bare grnd	25	
Lyonia luc	5 Bare grnd	75	
Ari str	60 Bare grnd	90	
Bare grnd	40 Bare grnd	70	
13 lle gla	60 Bare grnd	85	1997
Bare grnd	30 Bare grnd	75	66.56667
14 Lyonia luc	10 Cli mon	5	
Cli mon	10 Cli mon	3	
Ari str	50 Cli mon	15	
Bare grnd	5 Cli mon	15	
15 Lyonia luc	95 Cli mon	20	
Bare grnd	10 Cli mon	15	
16 Ari str	90 Cli mon	10	
Bare grnd	25 Cli mon	10	
17 Ari str			

Rhyn mic	5 Ci mon	10	
Ci mon	10 Ci mon	5	
Bare grnd	60 Ci mon	20	138
18 Ari str	5 Ci mon	10	4.6
rhyn mic	5 cyr rac	10	
Ci mon	5 cyr rac	20	-1
Bare grnd	85 lle gla	40	1.333333
19 Rhyn mic	5 llex mryt	65	2.166667
Bare grnd	95 lach car	5	0.166667
20 Rhyn mic	5 Lyonia luc	5	
Ari str	10 Lyonia luc	10	
Bare grnd	85 Lyonia luc	15	
21 llex mryt	65 Lyonia luc	20	
Lyonia luc	10 Lyonia luc	30	
Lyonia luc	25 Lyonia luc	20	
22 Ci mon	20 Lyonia luc	35	
Ari str	10 Lyonia luc	20	
Bare grnd	70 Lyonia luc	15	
23 Lyonia luc	25 Lyonia luc	25	
Ari str	10 Lyonia luc	20	
Bare grnd	65 Lyonia luc	25	
24 lach car	5 Lyonia luc	30	
Rhyn mic	65 Lyonia luc	5	
Ci mon	10 Lyonia luc	10	
Bare grnd	20 Lyonia luc	25	
25 Lyonia luc	75 Lyonia luc	25	
Bare grnd	25 Lyonia luc	75	
26 Lyonia luc	25 Lyonia luc	25	
Bare grnd	75 Lyonia luc	5	
27 Lyonia luc	5 Lyonia luc	15	
Rhyn mic	5 Lyonia luc	5	485
Bare grnd	90 Lyonia luc	25	16.16667
28 Ari str	10 Os amer	5	0.166667
Lyonia luc	15 Per pal	20	0.666667

Rhyn mic	5 Rhyn mic
Bare grnd	70 rhyn mic
29 Lyonia luc	5 Rhyn mic
Ari str	5 Rhyn mic
Vac cor	5 Rhyn mic
Bare grnd	85 Rhyn mic
30 Lyonia luc	25 Rhyn mic
Bare grnd	75 Vac cor

3000

5	5
5	5
5	5
5	5
65	95
5	3.166667
5	3.166667
5	0.167

Transect 7

Quadrat	Species
1	lach car
	Ari str
	Gay dum
	And glo
2	Bare ground
	Clif mon
	Ari str
	lach car
	Bare ground
3	Clif mon
	Ari str
	lach car
	Bare ground
4	Ari str
	And glo
	lach car
	Gay dum

Cover	Species
15	And glo
3	And glo
2	And glo
5	Ari str
75	Ari str
10	Ari str
3	Ari str
7	Ari str
80	Ari str
12	Ari str
5	Ari str
3	Ari str
80	Ari str
5	Ari str
5	Ari str
10	Ari str
10	Ari str

Cover

5	5
5	5
10	0.666667
3	3
3	3
5	5
5	5
15	15
5	5
5	5
5	5
3	3
10	10
5	5
5	5
5	5
5	5
84	84

- Bare ground
- Black titi
- Red root
- Red titi
- Wire grass
- Fetter bush
- Highbush blueberry
- Dwarf blueberry
- Bushy bluestem
- Dog fennel
- Gall berry
- Swamp dog hobbie
- Swamp bay
- Bunched beaksedge
- Myrtle-leaved holly

61.1
18.3
5.5
5.1
2.8
1.7
1.5
0.8
0.67
0.5
0.5
0.167
0.167
0.167
0.16

Species	Percent Cover
Bare ground	61.1
Black titi	18.3
Red root	5.5
Red titi	5.1
Mis species, ,8 (< 1% cover)	3.1
Wire grass	2.8
Fetter bush	1.7
Highbush blueberry	1.5
Species	Percent Cover
Dwarf blueberry	0.8
Bushy bluestem	0.67
Dog fennel	0.5
Gall berry	0.5
Swamp dog hobble	0.167
Swamp bay	0.167
Bunched beaksedge	0.167
Myrtle-leaved holly	0.16
Species	Percent Cover
70 Ari str	5
15 Bare ground	75
5 Bare ground	80
5 Bare ground	80
75 Bare ground	70
30 Bare ground	75
70 Bare ground	55
35 Bare ground	50
5 Bare ground	55
5 Bare ground	40
55 Bare ground	20
40 Bare ground	70
5 Bare ground	65
5 Bare ground	90
50 Bare ground	70
45 Bare ground	75
55 Bare ground	50
60 Bare ground	75
40 Bare ground	80
80 Bare ground	95
20 Bare ground	83
10 Bare ground	95
Rhyn mic	95
lach car	25
Clif mon	20
Lyon luc	45
Bare ground	70
13 Eup cap	35
Ari str	35
Clif mon	60
Bare ground	10
14 Lue rac	12
Clif mon	30
Bare ground	70
1833	61.1
3.131	

15 Pers pal	5 Clif mon	5	
Lyon luc	20 Clif mon	45	
Ari str	5 Clif mon	60	
Bare ground	70 Clif mon	80	
16 Ari str	3 Clif mon	5	
Gay dum	2 Clif mon	25	
Gay dum	10 Clif mon	5	
Lyon luc	10 Clif mon	25	
Bare ground	75 Clif mon	5	
17 Clif mon	25 Clif mon	10	
Ari str	10 Clif mon	2	
lach car	10 Clif mon	50	
Leuc rac	5 Clif mon	20	
Bare ground	50 Clif mon	25	
18 Clif mon	5 Clif mon	40	549
lach car	10 Clif mon	25	18.3
Lyon luc	5 Cyr rac	50	
Ari str	5 Cyr rac	45	
Bare ground	75 Cyr rac	60	5.166667
19 Ilex cor	15 Eup cap	5	
Ari str	5 Eup cap	10	0.5
Bare ground	80 Gay dum	2	
20 Lyon luc	5 Gay dum	10	
Bare ground	95 Gay dum	2	
21 Ari str	5 Gay dum	10	0.8
Lyon luc	2 Ilex cor	15	0.5
Clif mon	10 Ilex myr	5	0.166667
Bare ground	83 lach car	15	
22 Ari str	5 lach car	7	
Bare ground	95 lach car	3	
23 Clif mon	2 lach car	10	
Lue rac	3 lach car	5	
Bare ground	95 lach car	35	
24 Clif mon	50 lach car	40	

Vac corm	25 lach car	5	
Bare ground	25 lach car	10	
25 Cyr rac	50 lach car	10	165
Clif mon	20 lach car	25	5.5
Lue rac	10 Leuc rac	5	
Bare ground	20 Lue rac	5	
26 Cyr rac	45 Lue rac	3	
Vac corm	10 Lue rac	10	
Bare ground	45 Lue rac	5	0.166667
27 Clif mon	25 Lyon luc	5	
Lue rac	5 Lyon luc	5	
Bare ground	70 Lyon luc	20	
28 lach car	25 Lyon luc	10	
Clif mon	40 Lyon luc	5	
Bare ground	35 Lyon luc	5	
29 Cyr rac	60 Lyon luc	2	1.733333
llex myr	5 Pers pal	5	0.166667
Bare ground	35 Rhyrn mic	5	0/167
30 Eup cap	10 Vac corm	5	
Ari str	5 Vac corm	5	
Clif mon	25 Vac corm	25	
Bare ground	60 Vac corm	10	1.5

3000

Quadrat	Species	Cover	Species	Cover	Scientific Name	Species
Transect 8						
1	Carex ver	15	And glo	5	Bare ground	

Lach car	5 Bare ground	70		Muscadine grape	9.5
Rub arg	5 Bare ground	60		Panic grass	8
Rhex alf	2 Bare ground	95		Red root	6.5
Pin pal	3 Bare ground	55		Misc sp. 16, (<1% cover)	5.2
Bare ground	70 Bare ground	45		Black titi	5.1
2 Myr cer	25 Bare ground	75		Witch grass	3.16
Ilex gla	10 Bare ground	10		Poor joe	2.3
Lach car	5 Bare ground	60		Wax myrtle	1.8
Bare ground	60 Bare ground	85		Dog fennel	1.5
3 Lach car	5 Bare ground	35		Sand black berry	1.3
Bare ground	95 Bare ground	50		Partridge berry	1.1
4 Lach car	45 Bare ground	20		St. John's wort	1
Bare ground	55 Bare ground	65		Yaupon	1
5 Diod tere	5 Bare ground	85			
Lach car	50 Bare ground	35		Centella	0.8
Bare ground	45 Bare ground	75		Swamp bay	0.8
6 Vac corym	10 Bare ground	85		Gall berry	0.5
Lach car	10 Bare ground	55		Seedbox	0.5
Pan sp.	5 Bare ground	25		Warty Sedge	0.5
Bare ground	75 Bare ground	30		Jessamine	0.33
7 Pan sp.	50 Bare ground	35		Highbush blueberry	0.3
Cliff car	10 Bare ground	25		Water horehound	0.3
Per pal	15 Bare ground	30		Bunched beaksedge	0.167
Lach car	10 Bare ground	20		Bushy bluestem	0.167
Pan sp.	5 Bare ground	35		Cat briar	0.167
Bare ground	10 Bare ground	65		Red titi	0.167
8 Lach car	10 Bare ground	70		Swamp dog hobble	0.167
Myr cer	15 Bare ground	45		Candy weed	0.16
Per pal	5 Bare ground	65	1555	Longleaf pine	0.1
	10 Bare ground	50	51.833333	Savannah meadow beauty	0.06
	60 Carex ver	15	0.5		
	5 Cent as	5			
	5 Cent as	20	0.83333333		5.185
	5 Cham nic	10			

Bare ground	85 Cham nic	20	
Muscadine grape	40 Cham nic	5	1.166667
Panic grass	5 Clif car	10	
Red root	20 Clif car	10	
Misc sp. 16, (< 1% cover)	35 Clif car	5	
Black titi	5 Clif car	40	
Witch grass	5 Clif car	45	
Poor joe	5 Clif car	5	
Wax myrtle	5 Clif car	20	
Dog fennel	25 Clif car	10	155
Sand black berry	50 Clif car	5	5.166667
Partridge berry	45 Cyr rac	5	0.166667
St. John's wort	10 Dich sp.	5	
Yaupon	5 Dich sp.	5	
	20 Dich sp.	5	
	20 Dich sp.	40	
	5 Dich sp.	10	
	10 Dich sp.	10	
	65 Dich sp.	20	95
	5 Diod tere	5	3.166667
14 Lact car	5 Diod tere	5	
Dich sp.	5 Diod tere	5	
Clif car	5 Diod tere	5	
Bare ground	85 Diod tere	5	
15 Vit rotund	5 Diod tere	5	
Diod tere	5 Diod tere	5	
Pan sp.	10 Diod tere	20	
Dich sp.	40 Diod tere	10	
Clif car	5 Diod tere	5	70
Bare ground	35 Diod tere	5	2.333333
16 Ilex vom	5 Eup cap	45	1.5
lach car	5 Gel sem	5	
Cent as	5 Gel sem	5	3.333333

Diod tere	5	Hyper sp	5	
Pan sp.	5	Hyper sp	5	
Bare ground	75	Hyper sp	20	1
17 Diod tere	5	Ilex gla	10	
Lach car	5	Ilex gla	5	0.5
rbyn mic	5	Ilex vom	5	
Bare ground	85	Ilex vom	25	1
18 Lud dec	5	Lach car	5	
Per pal	20	Lach car	5	
Diod tere	5	Lach car	5	
Dich sp.	10	Lach car	45	
Hyper sp	5	Lach car	50	
Bare ground	55	Lach car	10	
19 Clif car	20	Lach car	10	
Cyr rac	5	Lach car	10	
Leuc rac	5	Lach car	5	
Hyper sp	20	Lach car	5	
Dich sp.	20	Lach car	5	
Diod tere	5	Lach car	10	
Bare ground	25	Lach car	5	
20 Cent as	20	Lach car	5	
Diod tere	20	Lach car	5	
Pan sp.	20	Lach car	5	6.5
Lud dec	5	Lach car	10	195
Lach car	5	Leuc rac	5	0.166667
Bare ground	30	Lud dec	5	
21 Ilex vom	25	Lud dec	5	0.5
Clif car	10	Lyc vir	10	0.333333
Lach car	10	Myr cer	25	
Pan sp.	20	Myr cer	15	
Bare ground	35	Myr cer	10	
22 Gel sem	5	Myr cer	5	1.833333
Pan sp.	40	Pan sp.	5	
Vit rotund	15	Pan sp.	50	

Diod tere	10 Pan sp.	5	
Rub arg	5 Pan sp.	20	
Bare ground	25 Pan sp.	25	
23 Lyc vir	10 Pan sp.	20	
Vit rotund	15 Pan sp.	20	
Eup cap	45 Pan sp.	10	
Bare ground	30 Pan sp.	5	
24 Vit rotund	60 Pan sp.	20	
Rub arg	10 Pan sp.	20	
Myr cer	10 Pan sp.	40	8
Bare ground	20 Per pal	15	
25 Vit rotund	50 Per pal	5	240
Rub arg	5 Per pal	20	0.833333
Cham nic	10 Pin pal	3	0.1
Bare ground	35 Poly lut	5	0.166667
26 Vit rotund	35 Rhex alf	2	0.066667
Bare ground	65 rhyn mic	5	0.166667
27 Vit rotund	25 Rub arg	5	
Dich sp.	5 Rub arg	5	
Bare ground	70 Rub arg	10	
28 Clif car	5 Rub arg	5	
Diod tere	5 Rub arg	10	
Vit rotund	25 Rub arg	5	1.333333
Cham nic	20 Smilax laur	5	0.167
Bare ground	45 Vac corym	10	0.333333
29 Smilax laur	5 Vit rotund	5	
Vit rotund	25 Vit rotund	15	
Myr cer	5 Vit rotund	15	
Bare ground	65 Vit rotund	60	
30 And glo	5 Vit rotund	50	
Vit rotund	30 Vit rotund	35	
Gel sem	5 Vit rotund	25	
Diod tere	5 Vit rotund	25	
Cham nic	5 Vit rotund	25	9.5

Bare ground

50 Vit rotund

30

285

3000

Quadrat # Transect 9

Species	Cover	Species	Cover	Common Name	Percent Cover
1 Rynch sp	40	Dros int	5	Open water	38
Dros int	5	Eleo cell	15	Fragrant water lily	35.5
Lach car	15	Eleo cell	15	Websteria	7.5
Open water	40	Eleo cell	20	Horned beaksedge	6.8
2 Nym od	50	Eleo cell	20	Spatterdock	4.7
Rynch inund	20	Eleo cell	20	Eleocharis	3.3
Open water	30	Eleo cell	10	Yellow-eyed grass	2.3
3 Eleo cell	15	Lach car	15	Florida bladderwort	1.2
Nym od	25	Nelumbo un	20	Mis species 2, (< 1 % co	0.66
Rynch inund	10	Nelumbo un	50	Red root	0.5
Open water	50	Nelumbo un	25	Water sundew (Fl threa	0.16
4 Nym od	15	Nelumbo un	40		4.5
	60	Nym od	50		
	25	Nym od	25		0.66
	35	Nym od	15		
	25	Nym od	35		
	40	Nym od	35		
	20	Nym od	15		
	35	Nym od	75		
	45	Nym od	15		
	60	Nym od	55		
	15	Nym od	50		
	25	Nym od	75		

■ Open water

■ Fragrant water lily

■ Websteria

- Horned beaksedge
- Spatterdock
- Eleocharis
- Yellow-eyed grass
- Florida bladderwort
- Mis species 2, (< 1 % cover)

Open water	10 Nym od	60	
13 Nym od	75 Nym od	50	1065
Open water	15 Nym od	55	
14 Nym od	15 Nym od	55	35.5
Utric flor	15 Nym od	40	
Open water	15 Nym od	65	
15 Nym od	55 Nym od	30	
Utric flor	40 Nym od	40	
Open water	60 Nym od	30	
16 Nym od	55 Nym od	40	
Open water	15 Nym od	45	
17 Nym od	30 Nym od	25	
Open water	50 Nym od	40	
18 Nym od	50 Nym od	20	
Eleo cell	75 Nym od	25	
Open water	25 Open water	40	
19 Nym od	60 Open water	30	
Open water	5 Open water	50	
20 Nym od	35 Open water	25	
Eleo cell	50 Open water	40	
Open water	20 Open water	45	
20 Nym od	40 Open water	35	
Open water	65 Open water	30	
Eleo cell	35 Open water	45	
Open water	30 Open water	45	
20 Nym od	20 Open water	40	
Open water	50 Open water	35	

21 Nym od	40 Open water	50	
Eleo cell	20 Open water	40	
Open water	40 Open water	30	
22 Nym od	30 Open water	50	
Webs conf	20 Open water	30	
Eleo cell	10 Open water	10	
Utric flor	10 Open water	45	
Open water	30 Open water	50	
23 Webs conf	10 Open water	20	
Nym od	40 Open water	60	1135
Open water	50 Open water	30	37.833333
24 Nelumbo un	20 Rhynch inund	10	
Rhynch inund	50 Rhynch inund	50	
Open water	30 Rhynch inund	40	
25 Nelumbo un	50 Rhynch inund	10	
Rhynch inund	40 Rhynch inund	25	
Open water	10 Rhynch inund	20	
26 Nym od	45 Rynch inund	20	
Rhynch inund	10 Rynch sp	40	6.8333333
Open water	45 Utric flor	5	
27 Nelumbo un	25 Utric flor	20	
Nym od	25 Utric flor	10	1.166667
Open water	50 Webs conf	60	
28 Nelumbo un	40 Webs conf	25	
Nym od	40 Webs conf	60	
Open water	20 Webs conf	10	
29 Nym od	20 Webs conf	20	
Webs conf	20 Webs conf	10	
Open water	60 Webs conf	20	
30 Rhynch inund	25 Webs conf	20	7.5
Nym od	25 Xyris sp	15	
Webs conf	20 Xyris sp	40	
Open water	30 Xyris sp	15	2.3333333

3000

3000